# abcam

## Product datasheet

## Anti-Myeloid Precursor Cells antibody [ER-MP58] ab63860

## 2 References

概述

产品名称 Anti-Myeloid Precursor Cells抗体[ER-MP58]

**描述** 大鼠单克隆抗体[ER-MP58] to Myeloid Precursor Cells

**宿主** Rat

经测试应用 适用于: Flow Cyt, IHC-Fr, ICC/IF, IP

种属反应性 与反应: Mouse

免疫原 Tissue, cells or virus corresponding to Mouse Myeloid Precursor Cells. Balb/c macrophage

precursor cell hybrids

常规说明

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

性能

形式 Liquid

**存放说明** Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term.

**存储溶液** pH: 7.40

Preservative: 0.09% Sodium azide

Constituent: PBS

纯**度** Protein G purified

**克隆** 单克隆

**克隆编号** ER-MP58 **骨髓瘤** Y3/Ag1.2.3

同种型 lgG2a

应用

1

#### The Abpromise guarantee

### Abpromise™承诺保证使用ab63860于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt		
IHC-Fr		
ICC/IF		
IP		

应**用说明** Flow Cyt: Use 10μl for 10<sup>6</sup> cells.

IHC-P: Use at an assay dependent dilution (PMID 19720836).

IHC-Fr: 1/25 - 1/100.

IP: Use at an assay dependent dilution.

Not yet tested in other applications.

Optimal dilutions/concentrations should be determined by the end user.

靶标

相关性 ER-MP58 is expressed by all bone marrow derived M-CSF- and GM-CSF-responsive myeloid

blood cell precursors. The expression of ER-MP58 remains at a high level throughout the

 $precursor/monocyte\ stage\ and\ is\ down\ regulated\ upon\ maturation\ into\ mature\ macrophages.\ The$ 

ER-MP58 antigen is used to distinguish between early myeloid committed cells and other

haematopoietic progenitors cells in the BM.

细**胞定位** Cell Membrane

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.cn/abpromise">https://www.abcam.cn/abpromise</a> or contact our technical team.

#### Terms and conditions

3

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors