abcam

Product datasheet

Anti-Myoglobin antibody [EP3081Y] ab77232



重组 RabMAb

★★★★★ 1 Abreviews 29 References 14 图像

概述

产品名称 Anti-Myoglobin抗体[EP3081Y]

描述 兔单克隆抗体[EP3081Y] to Myoglobin

宿主 Rabbit

经测试应用 适用于: WB, IHC-P, ICC/IF 种属反应性 与反应: Mouse, Rat, Human

免疫原 Synthetic peptide within Human Myoglobin aa 100-200 (C terminal). The exact sequence is

proprietary.

Database link: P02144

阳性对照 Human, rat and mouse muscle tissue lysates; human heart muscle tissue.

常规说明 This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity

- Long-term security of supply

- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.


性能

形式 Liquid

存放说明 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C.

Avoid freeze / thaw cycle.

 $K_D = 1.78 \times 10^{-10} M$ 解离常数(K_□)



Learn more about K_D

存储溶液 pH: 7.20 Preservative: 0.01% Sodium azide

Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.5% BSA

纯**度** Protein A purified

克隆编号 EP3081Y

同种型 IgG

应用

The Abpromise guarantee Abpromise™承诺保证使用ab77232于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。


应用	Ab评论	说明
WB	*****(1)	1/1000 - 1/5000. Predicted molecular weight: 17 kDa.
IHC-P		1/250 - 1/500. Perform heat mediated antigen retrieval via the pressure cooker method before commencing with IHC staining protocol. Use of an HRP/AP polymerized secondary antibody is recommended. See IHC antigen retrieval protocols.
ICC/IF		Use at an assay dependent concentration.

靶标


功能 Serves as a reserve supply of oxygen and facilitates the movement of oxygen within muscles.

序列相似性 Belongs to the globin family.

图片



Western blot - Anti-Myoglobin antibody [EP3081Y] (ab77232)



Lane 1: Human muscle lysate with 5% NFDM/TBST

Lane 2: Mouse muscle lysate with 5% NFDM/TBST

Lane 3: Rat muscle lysate with 5% NFDM/TBST

Lysates/proteins at 20 µg per lane.


Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (<u>ab97051</u>) at 1/20000 dilution (Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated)


Predicted band size: 17 kDa **Observed band size:** 17 kDa

Exposure time: 1 second

Purified ab77232 staining Myoglobin in Human cardiac muscle tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with paraformaldehyde; antigen retrieval was by heat mediation in a Tris/EDTA buffer (ab93684 Tris/EDTA buffer, pH 9.0). Samples were incubated with primary antibody (1/1000). A HRP-conjugated Goat Anti-Rabbit IgG H&L (ready to use) was used as the secondary antibody. Counterstained with Hematoxylin.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Myoglobin antibody
[EP3081Y] (ab77232)




Immunocytochemistry/ Immunofluorescence - Anti-Myoglobin antibody [EP3081Y] (ab77232)

Purified ab77232 staining Myoglobin in the C2C12 (mouse myoblasts myoblast) cell line by ICC/IF (Immunocytochemistry/immunofluorescence). Cells were fixed with methanol. Samples were incubated with primary antibody and an Alexa Fluor[®] 488-conjugated Goat anti-rabbit IgG (ab150077) was used as the secondary antibody. Counterstained with ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor[®] 594) and DAPI.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Myoglobin antibody
[EP3081Y] (ab77232)

Purified ab77232 staining Myoglobin in Mouse cardiac muscle tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with paraformaldehyde; antigen retrieval was by heat mediation in a Tris/EDTA buffer (ab93684 Tris/EDTA buffer, pH 9.0). Samples were incubated with primary antibody (1/1000). A HRP-conjugated Goat Anti-Rabbit IgG H&L (ready to use) was used as the secondary antibody. Counterstained with Hematoxylin.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Myoglobin antibody
[EP3081Y] (ab77232)


Purified ab77232 staining Myoglobin in Rat cardiac muscle tissue sections by Immunohistochemistry (IHC-P - paraformaldehydefixed, paraffin-embedded sections). Tissue was fixed with paraformaldehyde; antigen retrieval was by heat mediation in a Tris/EDTA buffer (ab93684 Tris/EDTA buffer, pH 9.0). Samples were incubated with primary antibody (1/1000). A HRP-conjugated Goat Anti-Rabbit IgG H&L (ready to use) was used as the secondary antibody. Counterstained with Hematoxylin.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Myoglobin antibody
[EP3081Y] (ab77232)

Unpurified ab77232 showing positive staining in human Skeletal muscle tissue.

Perform heat mediated antigen retrieval via the pressure cooker method before commencing with IHC staining protocol.




in 1x PBS + 0.1% Tween-20 for 17 hours at 4°C. A IRDye[®] 680-conjugated Donkey anti-rabbit polyclonal (1/40000) was used as the secondary antibody.

Western blot analysis of Pig skeletal muscle whole tissue lysate (10µg/lane) labelling Myoglobin with unpurified ab77232 at 1/5000

Western blot - Anti-Myoglobin antibody [EP3081Y] (ab77232)

This image is courtesy of an anonymous Abreview



Western blot - Anti-Myoglobin antibody [EP3081Y] (ab77232)

All lanes : Anti-Myoglobin antibody [EP3081Y] (ab77232) at 1/5000 dilution (Unpurified)

Lane 1 : Human heart muscle lysate
Lane 2 : Mouse heart muscle lysate

Lane 3: Rat heart muscle lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes: HRP labelled goat anti-rabbit at 1/2000 dilution


Predicted band size: 17 kDa Observed band size: 17 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Myoglobin antibody
[EP3081Y] (ab77232)

Unpurified ab77232, at a 1/250 dilution, staining Myoglobin in paraffin embedded human heart muscle by Immunohistochemistry.


Perform heat mediated antigen retrieval via the pressure cooker method before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Myoglobin antibody
[EP3081Y] (ab77232)

Unpurified ab77232 showing negative staining in Normal human brain tissue.


Perform heat mediated antigen retrieval via the pressure cooker method before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Myoglobin antibody
[EP3081Y] (ab77232)

Unpurified ab77232 showing negative staining in Normal human kidney tissue.

Perform heat mediated antigen retrieval via the pressure cooker method before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Myoglobin antibody
[EP3081Y] (ab77232)

Unpurified ab77232 showing negative staining in Normal human liver tissue.

Perform heat mediated antigen retrieval via the pressure cooker method before commencing with IHC staining protocol. Ol-RD Scanning - Anti-Myoglobin antibody [EP3081Y] (ab77232) Produced using unpurified ab77232.

Equilibrium disassociation constant (K_D)

Learn more about K_D

Click here to learn more about K_D



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors