

Anti-NAT10 antibody [EPR18663] ab194297

重组 RabMAb

★★★★★ **2 Abreviews** **11 References** **10 图像**

概述

产品名称	Anti-NAT10抗体[EPR18663]
描述	兔单克隆抗体[EPR18663] to NAT10
宿主	Rabbit
经测试应用	适用于: Flow Cyt (Intra), WB, ICC/IF, IP, IHC-P
种属反应性	与反应: Mouse, Rat, Human
免疫原	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: Human fetal brain and fetal heart lysates; Mouse brain, rat heart and rat spleen lysates. IHC-P: Human colon, mouse stomach and rat colon tissues. ICC/IF: HeLa and NIH/3T3 cells. IP: HeLa cell lysate. Flow Cyt (intra): NIH/3T3 cell lysate
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
存储溶液	<p>pH: 7.2</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA</p>
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR18663

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab194297于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt (Intra)		1/600.
WB	★★★★★ (1)	1/2000. Detects a band of approximately 116 kDa (predicted molecular weight: 116 kDa).
ICC/IF	★★★★★ (1)	1/2000.
IP		1/80.
IHC-P		1/500. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

靶标

功能

Has protein acetyltransferase activity in vitro. Can acetylate both histones and microtubules. Histone acetylation may regulate transcription and mitotic chromosome de-condensation. Activates telomerase activity by stimulating the transcription of TERT, and may also regulate telomerase function by affecting the balance of telomerase subunit assembly, disassembly, and localization. Acetylates alpha-tubulin, which may affect microtubule stability and cell division.


序列相似性

Belongs to the UPF0202 family.
Contains 1 N-acetyltransferase domain.

细胞定位

Nucleus > nucleolus. Nucleolar in interphase and redistributes to the perichromosomal layer and to the midbody during telophase.

图片



Immunocytochemistry/ Immunofluorescence - Anti-NAT10 antibody [EPR18663] (ab194297)

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HeLa (Human epithelial cells from cervix adenocarcinoma) cells labeling NAT10 with ab194297 at 1/2000 dilution, followed by Goat anti-rabbit IgG (Alexa Fluor® 488) (**ab150077**) secondary antibody at 1/1000 dilution (green).

Confocal image showing nuclear staining on HeLa cell line.


The nuclear counterstain is DAPI (blue).

Tubulin is detected with **ab7291** (anti-Tubulin mouse mAb) at 1/1000 dilution and **ab150120** (AlexaFluor®594 Goat anti-Mouse secondary) at 1/1000 dilution (red).

The negative controls are as follows:-

-ve control 1: ab194297 at 1/2000 dilution followed by **ab150120** (AlexaFluor®594 Goat anti-Mouse secondary) at 1/1000 dilution.

-ve control 2: **ab7291** (anti-Tubulin mouse mAb) at 1/1000 dilution followed by **ab150077** (Alexa Fluor®488 Goat Anti-Rabbit IgG H&L) at 1/1000 dilution.



Western blot - Anti-NAT10 antibody [EPR18663] (ab194297)

All lanes : Anti-NAT10 antibody [EPR18663] (ab194297) at 1/2000 dilution

Lane 1 : Mouse brain lysate

Lane 2 : Rat heart lysate

Lane 3 : Rat spleen lysate

Lysates/proteins at 10 µg per lane.


Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/50000 dilution

Predicted band size: 116 kDa

Observed band size: 116 kDa

Exposure time: 30 seconds



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-NAT10 antibody [EPR18663] (ab194297)


Immunohistochemical analysis of paraffin-embedded Rat colon tissue labeling NAT10 with ab194297 at 1/500 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution.

Nucleus staining on rat colon tissue is observed.

Counter stained with Hematoxylin.


Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-NAT10 antibody [EPR18663] (ab194297)

Intracellular Flow Cytometry analysis of NIH/3T3 (mouse embryo) cells labelling NAT10 (red) with purified ab194297 at dilution of 1/600. The secondary antibody used was Alexa Fluor[®] 488 goat-anti-rabbit IgG (1/2000). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. Isotype control antibody used was Rabbit Monoclonal IgG (black). The blue line shows cells without incubation with primary antibody and secondary antibody.



Western blot - Anti-NAT10 antibody [EPR18663] (ab194297)

All lanes : Anti-NAT10 antibody [EPR18663] (ab194297) at 1/2000 dilution

Lane 1 : Human fetal brain lysate

Lane 2 : Human fetal heart lysate

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG at 1/10000 dilution

Predicted band size: 116 kDa

Observed band size: 116 kDa

Exposure time: 30 seconds

Blocking/Dilution buffer: 5% NFDM/TBST.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-NAT10 antibody [EPR18663] (ab194297)


Immunohistochemical analysis of paraffin-embedded Human colon tissue labeling NAT10 with ab194297 at 1/500 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/500 dilution.

Nucleus staining on Human colon tissue is observed.

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/500 dilution.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunocytochemistry/ Immunofluorescence - Anti-NAT10 antibody [EPR18663] (ab194297)

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized NIH/3T3 (Mouse embryo fibroblast cells) cells labeling NAT10 with ab194297 at 1/2000 dilution, followed by Goat anti-rabbit IgG (Alexa Fluor® 488) (**ab150077**) secondary antibody at 1/1000 dilution (green).

Confocal image showing nuclear staining on NIH/3T3 cell line.


The nuclear counterstain is DAPI (blue).

Tubulin is detected with **ab7291** (anti-Tubulin mouse mAb) at 1/1000 dilution and **ab150120** (AlexaFluor®594 Goat anti-Mouse secondary) at 1/1000 dilution (red).

The negative controls are as follows:-

-ve control 1: ab194297 at 1/2000 dilution followed by **ab150120** (AlexaFluor®594 Goat anti-Mouse secondary) at 1/1000 dilution.

-ve control 2: **ab7291** (anti-Tubulin mouse mAb) at 1/1000 dilution followed by **ab150077** (Alexa Fluor®488 Goat Anti-Rabbit IgG H&L) at 1/1000 dilution.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-NAT10 antibody [EPR18663] (ab194297)


Immunohistochemical analysis of paraffin-embedded Mouse stomach tissue labeling NAT10 with ab194297 at 1/500 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/500 dilution.

Nucleus staining on mouse stomach tissue is observed.

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/500 dilution.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



NAT10 was immunoprecipitated from 1mg of HeLa (Human epithelial cells from cervix adenocarcinoma) cell lysate with ab194297 at 1/80 dilution.

Lane 1: HeLa cell lysate 10ug (Input).

Lane 2: ab194297 IP in HeLa cell lysate.

Lane 3: Rabbit monoclonal IgG (**ab172730**) instead of ab194297 in HeLa cell lysate.

Western blot was performed from the immunoprecipitate using ab194297 at 1/1000 dilution. Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG, was used as secondary antibody at 1/1500.

Blocking and dilution buffer and concentration: 5% NFD/MTBST.

Exposure time: 3 seconds.

Why choose a recombinant antibody?

<p>Research with confidence Consistent and reproducible results</p>	<p>Long-term and scalable supply Recombinant technology</p>
<p>Success from the first experiment Confirmed specificity</p>	<p>Ethical standards compliant Animal-free production</p>

Anti-NAT10 antibody [EPR18663] (ab194297)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors