

Anti-Neisseria gonorrhoeae antibody ab19962

2 References

概述

产品名称	Anti-Neisseria gonorrhoeae抗体
描述	兔多克隆抗体to Neisseria gonorrhoeae
宿主	Rabbit
特异性	Reacts to all antigens of Neisseria gonorrhoeae. Antiserum is not absorbed and may react with related microorganisms.
经测试应用	适用于: ICC/IF
种属反应性	与反应: Neisseria gonorrhoeae
免疫原	Tissue, cells or virus corresponding to Human Neisseria gonorrhoeae. Whole Neisseria gonorrhoeae.ATCC 31426
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	<p>pH: 7.20</p> <p>Preservative: 0.1% Sodium azide</p> <p>Constituent: 0.0268% PBS</p>
纯度	Protein A purified
克隆	多克隆
同种型	IgG

应用

The Abpromise guarantee

Abpromise™ 承诺保证使用ab19962于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC/IF		Use at an assay dependent concentration.

靶标

相关性	Neisseria gonorrhoeae infections are acquired by sexual contact and usually affect the mucous membranes of the urethra in males and the endocervix and urethra in females, although the infection may disseminate to a variety of tissues. The pathogenic mechanism involves the attachment of the bacterium to nonciliated epithelial cells via pili (fimbriae) and the production of lipopolysaccharide endotoxin. Similarly, the lipopolysaccharide of Neisseria meningitidis is highly toxic, as it has an additional virulence factor in the form of its antiphagocytic capsule. Both pathogens produce IgA proteases which promote virulence. Many normal individuals may harbor Neisseria meningitidis in the upper respiratory tract, but Neisseria gonorrhoeae is never part of the normal flora and is only found after sexual contact with an infected person (or direct contact, in the case of infections in the newborn).
-----	---

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors