

Anti-Netrin 1 antibody ab122903

[4 References](#) [2 图像](#)

概述

产品名称	Anti-Netrin 1抗体
描述	山羊多克隆抗体to Netrin 1
宿主	Goat
经测试应用	适用于: WB, IHC-P
种属反应性	与反应: Rat, Human 预测可用于: Cow, Dog 
免疫原	Synthetic peptide conjugated to KLH, corresponding to C terminal amino acids of Human Netrin 1.
阳性对照	Human Intestine tissue; Rat, Mouse and Human Brain lysates.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
存储溶液	pH: 7.30 Preservative: 0.02% Sodium azide Constituents: 99% Tris buffered saline, 0.5% BSA
纯度	Protein G purified
克隆	多克隆
同种型	IgG

应用

The Abpromise guarantee

Abpromise™承诺保证使用ab122903于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		Use a concentration of 0.5 - 1.5 µg/ml. Detects a band of approximately 70 kDa (predicted molecular weight: 68 kDa).
IHC-P		Use a concentration of 3.75 µg/ml.


靶标

功能	Netrins control guidance of CNS commissural axons and peripheral motor axons. Its association with either DCC or some UNC5 receptors will lead to axon attraction or repulsion, respectively. It also serve as a survival factor via its association with its receptors which prevent the initiation of apoptosis. Involved in tumorigenesis by regulating apoptosis.
组织特异性	Widely expressed in normal adult tissues with highest levels in heart, small intestine, colon, liver and prostate. Reduced expression in brain tumors and neuroblastomas.
序列相似性	Contains 3 laminin EGF-like domains. Contains 1 laminin N-terminal domain. Contains 1 NTR domain.
细胞定位	Secreted > extracellular space > extracellular matrix.

图片



ab122903, at 3.75 µg/ml, staining Netrin 1 in Formalin-fixed, Paraffin-embedded Human Intestine tissue by Immunohistochemistry.



Anti-Netrin 1 antibody (ab122903) at 0.5 µg/ml + Rat Brain lysate (in RIPA buffer) at 35 µg

Developed using the ECL technique.

Predicted band size: 68 kDa

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors