# abcam

# Product datasheet

# Anti-NEURL1B antibody ab122400

★★★★★ 1 Abreviews 3 图像

#### 概述

产**品名称** Anti-NEURL1B**抗体** 

描述 兔多克隆抗体to NEURL1B

**宿主** Rabbit

经测试应用 适用于: ICC/IF, IHC-P, WB

种属反应性 与反应: Human

免疫原 antigen: ALIDVYGITD EVQLLESAFA DTLTPARLSQ ARFSACLPPS SHDAANFDNN

ELENNQVVAK LGHL, corresponding to amino acids 178-241 of Human NEURL1B according to

A8MQ27

Run BLAST with EXPASY Run BLAST with S NCBI

**阳性**对照 Human rectum tissue; Human liver, Human tonsil, RT-4 and U-251 MG cell lysates.

常规说明

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

性能

形式 Liquid

**存放说明** Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

存储溶液 pH: 7.20

Preservative: 0.02% Sodium azide

Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine)

纯**度** Immunogen affinity purified

**克隆** 多克隆

同种型 lgG

应用

1

### The Abpromise guarantee

### Abpromise™承诺保证使用ab122400于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC/IF		Use a concentration of 0.25 - 2 µg/ml.
IHC-P		1/200 - 1/500. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
WB		Use a concentration of 0.04 - 0.4 μg/ml.

#### 靶标


功能 E3 ubiquitin-protein ligase.

通路 Protein modification; protein ubiquitination.

序列相似性 Contains 2 NHR (neuralized homology repeat) domains.


Contains 1 RING-type zinc finger.

### 图片




Immunocytochemistry/ Immunofluorescence - Anti-NEURL1B antibody (ab122400)

Immunofluorescent staining of Human cell line U-251 MG shows positivity in vesicles. Recommended concentration of ab122400 1-4  $\mu$ g/ml. Cells treated with PFA/Triton X-100.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-NEURL1B antibody (ab122400)

ab122400, at a 1/225 dilution, staining NEURL1B in glandular cells of paraffin-embedded Human rectum tissue by immunohistochemistry.



All lanes: Anti-NEURL1B antibody (ab122400) at 1/250 dilution

Lane 1: RT-4 cell lysate

Lane 2: U-251 MG cell lysate

Lane 3: Human plasma

Lane 4: Human liver lysate

Lane 5: Human tonsil lysate

Developed using the ECL technique.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.cn/abpromise">https://www.abcam.cn/abpromise</a> or contact our technical team.

#### Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors