

Anti-Nurr1 antibody [N1404] ab41917

★★★★★ [9 Abreviews](#) [24 References](#) [1 图像](#)

概述

产品名称	Anti-Nurr1抗体[N1404]
描述	小鼠单克隆抗体[N1404] to Nurr1
宿主	Mouse
经测试应用	适用于: IHC-P
种属反应性	与反应: Rat
免疫原	Recombinant fragment: CVQAQYGSSP QGASPASQSY SYHSSGEYSS DFLTPEFVKF SMDLTNTEIT ATTSLPSTFST FMDNYSTGYD VKPPCLYQMP LSGQQSSIKV EDIQMHN, corresponding to amino acids 2-99 of Human Nurr1 Run BLAST with Expasy Run BLAST with NCBI
阳性对照	Rat cerebral nerve cell.
常规说明	<p>This product was changed from ascites to tissue culture supernatant on 3rd April 2019. Please note that the dilutions may need to be adjusted accordingly. If you have any questions, please do not hesitate to contact our scientific support team.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
存储溶液	pH: 7 Preservative: 0.1% Sodium azide
纯度	Physiological saline. Tissue culture supernatant

克隆	单克隆
克隆编号	N1404
同种型	IgG2a

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab41917于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-P	★★★★★ (6)	Use at an assay dependent concentration.

靶标

功能	Transcriptional regulator which is important for the differentiation and maintenance of meso-diencephalic dopaminergic (mdDA) neurons during development. It is crucial for expression of a set of genes such as SLC6A3, SLC18A2, TH and DRD2 which are essential for development of mdDA neurons.
组织特异性	Expressed in a number of cell lines of T-cell, B-cell and fibroblast origin. Strong expression in brain tissue.
序列相似性	Belongs to the nuclear hormone receptor family. NR4 subfamily. Contains 1 nuclear receptor DNA-binding domain.
发展阶段	Rapidly and only very transiently expressed after cell activation, during the G0-G1 transition of the cell cycle.
细胞定位	Nucleus.

图片



ab41917 staining Nurr1 rat cerebral nerve cell paraffin section (10 ug/mL) by immunohistochemistry, formalin-fixed paraffin embedded sections.

This image was generated using the ascites version of the product.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Nurr1 antibody [N1404] (ab41917)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors