

Anti-p53R2 antibody [EPR8816] ab154194

敲除验证 重组 RabMAb

[6 References](#) [18 图像](#)

概述

产品名称	Anti-p53R2抗体[EPR8816]
描述	兔单克隆抗体[EPR8816] to p53R2
宿主	Rabbit
经测试应用	适用于: WB, IHC-P, ICC/IF, IP, Flow Cyt (Intra)
种属反应性	与反应: Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: MCF7, HCT116, Human skeletal muscle, and SW480 lysates. IHC-P: human breast carcinoma tissue. ICC/IF: HeLa cells. Flow Cyt (intra): MCF7 cells. IP: MCF7 cells.
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p> <p>Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at -20°C.
存储溶液	Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR8816
同种型	IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab154194于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/1000 - 1/10000. Predicted molecular weight: 40 kDa.
IHC-P		1/500. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. For unpurified use at 1/50 - 1/100
ICC/IF		1/50.
IP		1/10 - 1/100.
Flow Cyt (Intra)		1/10 - 1/100. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

靶标

功能	Plays a pivotal role in cell survival by repairing damaged DNA in a p53/TP53-dependent manner. Supplies deoxyribonucleotides for DNA repair in cells arrested at G1 or G2. Contains an iron-tyrosyl free radical center required for catalysis. Forms an active ribonucleotide reductase (RNR) complex with RRM1 which is expressed both in resting and proliferating cells in response to DNA damage.
组织特异性	Widely expressed at a high level in skeletal muscle and at a weak level in thymus. Expressed in epithelial dysplasias and squamous cell carcinoma.
通路	Genetic information processing; DNA replication.
疾病相关	Defects in RRM2B are the cause of mitochondrial DNA depletion syndrome type 8A (MTDPS8A) [MIM:612075]. A disorder due to mitochondrial dysfunction characterized by various combinations of neonatal hypotonia, neurological deterioration, respiratory distress, lactic acidosis, and renal tubulopathy. Defects in RRM2B are the cause of mitochondrial DNA depletion syndrome type 8B (MTDPS8B) [MIM:612075]. A disease due to mitochondrial dysfunction and characterized by ophthalmoplegia, ptosis, gastrointestinal dysmotility, cachexia, peripheral neuropathy. Defects in RRM2B are the cause of progressive external ophthalmoplegia with mitochondrial DNA deletions autosomal dominant type 5 (PEOA5) [MIM:613077]. A disorder characterized by progressive weakness of ocular muscles and levator muscle of the upper eyelid. In a minority of cases, it is associated with skeletal myopathy, which predominantly involves axial or proximal muscles and which causes abnormal fatigability and even permanent muscle weakness. Ragged-red fibers and atrophy are found on muscle biopsy. A large proportion of chronic ophthalmoplegias are associated with other symptoms, leading to a multisystemic pattern of this disease. Additional symptoms are variable, and may include cataracts, hearing loss, sensory axonal neuropathy, ataxia, depression, hypogonadism, and parkinsonism.


序列相似性

Belongs to the ribonucleoside diphosphate reductase small chain family.

细胞定位

Cytoplasm. Nucleus. Translocates from cytoplasm to nucleus in response to DNA damage.

图片



All lanes : Anti-p53R2 antibody [EPR8816] (ab154194) at 1/1000 dilution

Lane 1 : Wild-type HeLa cell lysate

Lane 2 : RRM2B knockout HeLa cell lysate

Lysates/proteins at 40 µg per lane.


Performed under reducing conditions.

Predicted band size: 40 kDa

Observed band size: 40 kDa


Lanes 1-2: Merged signal (red and green). Green - ab154194 observed at 40 kDa. Red - Anti-alpha Tubulin antibody [DM1A] - Loading Control ([ab7291](#)) observed at 50 kDa.

ab154194 was shown to react with p53R2 in wild-type HeLa cells in western blot. Loss of signal was observed when knockout cell line [ab261769](#) (knockout cell lysate [ab257215](#)) was used. Wild-type HeLa and RRM2B knockout HeLa cell lysates were subjected to SDS-PAGE. Membrane was blocked for 1 hour at room temperature in 0.1% TBST with 3% non-fat dried milk. ab154194 and Anti-alpha Tubulin antibody [DM1A] - Loading Control ([ab7291](#)) overnight at 4°C at a 1 in 1000 dilution and a 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye®800CW) preadsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye®680RD) preadsorbed ([ab216776](#)) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.




Immunocytochemistry/ Immunofluorescence - Anti-p53R2 antibody [EPR8816] (ab154194)

Immunocytochemistry/ Immunofluorescence analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling p53R2 with purified ab154194 at 1/50 dilution (2.2 µg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with **ab195889** Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1/200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) was used as the secondary antibody at 1/1000 (2 µg/ml) dilution. DAPI (blue) was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-p53R2 antibody [EPR8816] (ab154194)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human breast carcinoma tissue sections labeling p53R2 with purified ab154194 at 1/500 dilution (0.22 µg/ml). Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 2 (pH 9.0). Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Flow Cytometry (Intracellular) - Anti-p53R2 antibody [EPR8816] (ab154194)

Intracellular Flow Cytometry analysis of MCF7 (Human breast adenocarcinoma epithelial cell) cells labeling p53R2 with purified ab154194 at 1/20 dilution (10µg/ml) (Red). Cells were fixed with 4% Paraformaldehyde and permeabilised with 90% Methanol. A Goat anti rabbit IgG (Alexa Fluor® 488, [ab150077](#)) secondary antibody was used at 1/2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).



Western blot - Anti-p53R2 antibody [EPR8816] (ab154194)

All lanes : Anti-p53R2 antibody [EPR8816] (ab154194) at 1/1000 dilution

Lane 1 : Wild-type HCT116 cell lysate

Lane 2 : RRM2B knockout HCT116 cell lysate

Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.


Predicted band size: 40 kDa

Observed band size: 40 kDa

Lanes 1- 2: Merged signal (red and green). Green - ab154194 observed at 40 kDa. Red - Anti-alpha Tubulin antibody [DM1A] - Loading Control ([ab7291](#)) observed at 50 kDa.

ab154194 was shown to react with p53R2 in wild-type HCT116 cells in western blot. Loss of signal was observed when knockout cell line [ab266897](#) (knockout cell lysate [ab257216](#)) was used. Wild-Type HCT116 and RRM2B knockout HCT116 cell lysates were

subjected to SDS-PAGE. Membrane was blocked for 1 hour at room temperature in 0.1% TBST with 3% non-fat dried milk. ab154194 and Anti-alpha Tubulin antibody [DM1A] - Loading Control ([ab7291](#)) overnight at 4°C at a 1 in 1000 dilution and a 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye®800CW) preadsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye®680RD) preadsorbed ([ab216776](#)) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Western blot - Anti-p53R2 antibody [EPR8816] (ab154194)

All lanes : Anti-p53R2 antibody [EPR8816] (ab154194) at 1/1000 dilution (Purified)

Lane 1 : MCF7 (Human breast adenocarcinoma epithelial cell) whole cell lysates

Lane 2 : Human skeletal muscle lysates

Lane 3 : SW480 (Human colorectal adenocarcinoma epithelial cell) whole cell lysates


Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG (HRP) with minimal cross-reactivity with human IgG at 1/2000 dilution

Predicted band size: 40 kDa

Observed band size: 41 kDa



Western blot - Anti-p53R2 antibody [EPR8816] (ab154194)

Lane 1 : Wild-type HAP1 cell lysate (20 µg)

Lane 2 : p53R2 knockout HAP1 cell lysate (20 µg)


Lane 3 : MCF7 cell lysate (20 µg)

Lane 4 : SW480 cell lysate (20 µg)

Lanes 1 - 4: Merged signal (red and green). Green - ab154194 observed at 40 kDa. Red - loading control, [ab18058](#), observed at 124 kDa.

Unpurified ab154194 was shown to recognize p53R2 when p53R2 knockout samples were used, along with additional cross-reactive bands. Wild-type and p53R2 knockout samples were subjected to SDS-PAGE. ab154194 and [ab18058](#) (loading control to Vinculin) were both diluted 1/1000 and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye®

680RD) preadsorbed (**ab216776**) secondary antibodies at 1/10000 dilution for 1 h at room temperature before imaging.



Western blot - Anti-p53R2 antibody [EPR8816] (ab154194)

All lanes : Anti-p53R2 antibody [EPR8816] (ab154194) at 1/1000 dilution ((unpurified))

Lane 1 : Human fetal muscle lysate

Lane 2 : MCF7 cell lysate


Lane 3 : SW480 cell lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat anti-rabbit HRP at 1/2000 dilution

Predicted band size: 40 kDa



Immunoprecipitation - Anti-p53R2 antibody [EPR8816] (ab154194)

ab154194 (purified) at 1/20 dilution (0.5ug) immunoprecipitating p53R2 in MCF7 whole cell lysate.


Lane 1 (input): MCF7 (Human breast adenocarcinoma epithelial cell) whole cell lysate 10ug

Lane 2 (+): ab154194 & MCF7 whole cell lysate

Lane 3 (-): Rabbit monoclonal IgG (**ab172730**) instead of ab154194 in MCF7 whole cell lysate


For western blotting, VeriBlot for IP Detection Reagent (HRP) (**ab131366**) was used at 1/1000 dilution.

Blocking and diluting buffer: 5% NFDN/TBST.




Immunocytochemistry/ Immunofluorescence - Anti-p53R2 antibody [EPR8816] (ab154194)

Immunofluorescent staining of MCF7 cells labeling p53R2 with unpurified ab154194 at 1/250 dilution.



Flow Cytometry (Intracellular) - Anti-p53R2 antibody
[EPR8816] (ab154194)


Intracellular flow cytometric analysis of permeabilized MCF7 cells labeling p53R2 with unpurified ab154194 at 1/10 dilution (red) or a rabbit IgG negative control antibody (green).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-p53R2 antibody
[EPR8816] (ab154194)

Immunohistochemical analysis of paraffin-embedded Human ovarian carcinoma tissue labeling p53R2 with unpurified ab154194 at 1/50 dilution.


Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-p53R2 antibody
[EPR8816] (ab154194)

Immunohistochemical analysis of paraffin embedded Human urinary bladder transitional carcinoma tissue using unpurified ab154194 showing +ve staining.


Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemical analysis of paraffin embedded Human melanoma tissue using unpurified ab154194 showing +ve staining.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-p53R2 antibody [EPR8816] (ab154194)



Immunohistochemical analysis of paraffin embedded Human normal kidney tissue unpurified ab154194 showing +ve staining.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-p53R2 antibody [EPR8816] (ab154194)



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-p53R2 antibody [EPR8816] (ab154194)

Immunohistochemical analysis of paraffin embedded Human normal colon tissue using unpurified ab154194 showing +ve staining.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-p53R2 antibody [EPR8816] (ab154194)


Immunohistochemical analysis of paraffin embedded Human normal brain tissue using unpurified ab154194 showing +ve staining.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.

Why choose a recombinant antibody?

- 

Research with confidence
Consistent and reproducible results
- 

Long-term and scalable supply
Recombinant technology
- 

Success from the first experiment
Confirmed specificity
- 

Ethical standards compliant
Animal-free production

Anti-p53R2 antibody [EPR8816] (ab154194)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors