

# Anti-p75 NGF Receptor antibody [MC-192] ab6172

★★★★★ [2 Abreviews](#) [13 References](#) [1 图像](#)

### 概述

产品名称	Anti-p75 NGF Receptor抗体[MC-192]
描述	小鼠单克隆抗体[MC-192] to p75 NGF Receptor
宿主	Mouse
经测试应用	适用于: ICC
种属反应性	与反应: Rat
免疫原	Tissue, cells or virus corresponding to Rat p75 NGF Receptor. PC-12 cells.
阳性对照	This antibody gave a positive result when used in the following formaldehyde fixed cell lines: PC12.
常规说明	<p>A well characterised and versatile monoclonal antibody, clone number MC-192, that is of central importance in the study of LANGF (p75-NGF) receptor.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Add glycerol to a final volume of 50% for extra stability and aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	pH: 7 Constituent: PBS
纯度	Protein A purified
Primary antibody说明	A well characterised and versatile monoclonal antibody, clone number MC-192, that is of central importance in the study of LANGF (p75-NGF) receptor.
克隆	单克隆

克隆编号	MC-192
同种型	IgG1

## 应用

**The Abpromise guarantee**      **Abpromise™** 承诺保证使用ab6172于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC		Use at an assay dependent concentration.

## 靶标

功能	Low affinity receptor which can bind to NGF, BDNF, NT-3, and NT-4. Can mediate cell survival as well as cell death of neural cells.
序列相似性	Contains 1 death domain. Contains 4 TNFR-Cys repeats.
结构域	Death domain is responsible for interaction with RANBP9. The extracellular domain is responsible for interaction with NTRK1.
翻译后修饰	N- and O-glycosylated. O-linked glycans consist of Gal(1-3)GalNAc core elongated by 1 or 2 NeuNAc. Phosphorylated on serine residues.
细胞定位	Membrane.

## 图片



Copyright (c) 2012 abcam plc

Immunocytochemistry - Anti-p75 NGF Receptor antibody [MC-192] (ab6172)

ab6172 stained PC12 cells. The cells were 4% formaldehyde fixed (10 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody ab6172 at 5µg/ml overnight at +4°C. The secondary antibody (green) was DyLight® 488 goat anti-mouse (**ab96879**) IgG (H+L) used at a 1/1000 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors