

Anti-PABP antibody ab21060

★★★★★ [12 Abreviews](#) [120 References](#) [7 图像](#)

概述

产品名称	Anti-PABP抗体
描述	兔多克隆抗体to PABP
宿主	Rabbit
特异性	From Jan 2024, QC testing of replenishment batches of this polyclonal changed. All tested and expected application and reactive species combinations are still covered by our Abcam product promise. However, we no longer test all applications. For more information on a specific batch, please contact our Scientific Support who will be happy to help. You may also be interested in our alternative recombinant antibody, ab312314 .
经测试应用	适用于: WB, ICC/IF
种属反应性	与反应: Mouse, Rat, Human 预测可用于: Cow, Pig 
免疫原	Synthetic peptide conjugated to KLH derived from within residues 600 to the C-terminus of Human PABP. 参阅Abcam的专有抗源政策 (Peptide available as ab23710 .)
阳性对照	ICC/IF: HeLa cells. WB: HeLa, NIH/3T3, MEF1 and PC12 whole cell lysate. Mouse brain, testis, spinal cord, ovary and pancreas tissue lysate. Brain (Rat) Tissue Lysate - normal tissue.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	pH: 7.40 Preservative: 0.02% Sodium azide Constituent: PBS

Batches of this product that have a concentration < 1mg/ml may have BSA added as a stabilising agent. If you would like information about the formulation of a specific lot, please contact our scientific support team who will be happy to help.

纯度 Immunogen affinity purified
克隆 多克隆
同种型 IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab21060于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB	★★★★★ (4)	Use a concentration of 1 µg/ml. Detects a band of approximately 75 kDa.
ICC/IF	★★★★★ (4)	Use a concentration of 1 - 5 µg/ml.

靶标

功能 Binds the poly(A) tail of mRNA, including that of its own transcript. May be involved in cytoplasmic regulatory processes of mRNA metabolism such as pre-mRNA splicing. Its function in translational initiation regulation can either be enhanced by PAIP1 or repressed by PAIP2. Can probably bind to cytoplasmic RNA sequences other than poly(A) in vivo. Involved in translationally coupled mRNA turnover. Implicated with other RNA-binding proteins in the cytoplasmic deadenylation/translational and decay interplay of the FOS mRNA mediated by the major coding-region determinant of instability (mCRD) domain. Involved in regulation of nonsense-mediated decay (NMD) of mRNAs containing premature stop codons; for the recognition of premature termination codons (PTC) and initiation of NMD a competitive interaction between UPF1 and PABPC1 with the ribosome-bound release factors is proposed. By binding to long poly(A) tails, may protect them from uridylation by ZCCHC6/ZCCHC11 and hence contribute to mRNA stability (PubMed:25480299). Positively regulates the replication of dengue virus (DENV) (PubMed:26735137).

组织特异性 Ubiquitous.


序列相似性 Belongs to the polyadenylate-binding protein type-1 family.
Contains 1 PABC domain.
Contains 4 RRM (RNA recognition motif) domains.

结构域 The RNA-binding domains RRM1 and RRM2 and the C-terminus (last 138 amino acids) regions interact with the PABPC1-interacting motif-1 (PAM1) and -2 (PAM2) of PAIP1, respectively. The RNA-binding domains RRM2 and RRM3 and the C-terminus (last 138 amino acids) regions interact with the PABPC1-interacting motif-1 (PAM1) and -2 (PAM2) of PAIP2, respectively.


翻译后修饰 Phosphorylated by MAPKAPK2.
Methylated by CARM1. Arg-493 is dimethylated, probably to asymmetric dimethylarginine.

细胞定位 Cytoplasm. Nucleus. Localized in cytoplasmic mRNP granules containing untranslated mRNAs. Shuttles between the cytoplasm and the nucleus.

图片



Immunocytochemistry/ Immunofluorescence - Anti-PABP antibody (ab21060)



Western blot - Anti-PABP antibody (ab21060)

ab21060 staining PABP in HeLa cells. The cells were fixed with 100% methanol (5 min), permeabilized with 0.1% PBS-Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at 4°C with ab21060 at 1 µg/ml and **ab7291**, Mouse monoclonal [DM1A] to alpha Tubulin - Loading Control. Cells were then incubated with **ab150081**, Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (Alexa Fluor® 488), pre-adsorbed at 1/1000 dilution (shown in green) and **ab150120**, Goat polyclonal Secondary Antibody to Mouse IgG - H&L (Alexa Fluor® 594), pre-adsorbed at 1/1000 dilution (shown in pseudocolour red). Nuclear DNA was labelled with DAPI (shown in blue).

Also suitable in cells fixed with 4% paraformaldehyde (10 min).

Image was acquired with a high-content analyser (Operetta CLS, Perkin Elmer) and a maximum intensity projection of confocal sections is shown.

Anti-PABP antibody (ab21060) at 1 µg/ml + HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate at 20 µg/ml

Secondary


Goat polyclonal to Rabbit IgG (Alexa Fluor® 680) at 1/10000 dilution

Performed under reducing conditions.

Predicted band size: 69 kDa

Observed band size: 75 kDa

ab21060 detects a band of ~ 75 kDa in HeLa whole cell lysate. A 75 kDa band is also seen in Jurkat and A431 whole cell lysates (data not shown). PABP is reported to have run at an apparent molecular mass of 73 kDa on SDS-PAGE. We believe the band we see at 75 kDa represents PABP.




Immunocytochemistry/ Immunofluorescence - Anti-PABP antibody (ab21060)

Sharma et al PLoS Pathog. 2017 Oct 30;13(10):e1006677. doi: 10.1371/journal.ppat.1006677. eCollection 2017 Oct. Fig 2. Reproduced under the Creative Commons license <https://creativecommons.org/publicdomain/zero/1.0/>

Viral ORF57 expression is required to suppress the formation of SG during KSHV lytic infection and under arsenite stress.

ORF57 alone is sufficient to inhibit SG formation. HeLa cells transfected with an ORF57-Flag (pVM7) expressing vector or an empty vector (control) for 24 h were induced with 0.5 mM arsenite for 30 min for SG formation. The cells were stained for ORF57 (green), SG-specific TIA-1 (red) and PABPC1 (white) (ab21060) by each corresponding antibody. The nuclei were counterstained with Hoechst stain. Scale bar = 10 μ m.



Western blot - Anti-PABP antibody (ab21060)

All lanes : Anti-PABP antibody (ab21060) at 1 μ g/ml

Lane 1 : NIH/3T3 whole cell lysate ([ab7179](#))

Lane 2 : MEF1 (Mouse embryonic fibroblast cell line) Whole Cell Lysate

Lane 3 : Brain (Mouse) Tissue Lysate

Lane 4 : Mouse pancreas tissue lysate - total protein ([ab29363](#))

Lane 5 : Testis (Mouse) Tissue Lysate - normal tissue

Lane 6 : Spinal Cord (Mouse) Tissue Lysate

Lane 7 : Ovary (Mouse) Tissue Lysate - normal tissue

Lane 8 : PC-12 (Rat adrenal pheochromocytoma cell line) Whole Cell Lysate

Lane 9 : Brain (Rat) Tissue Lysate - normal tissue

Lysates/proteins at 10 μ g per lane.

Secondary


All lanes : IRDye 680 Conjugated Goat Anti-Rabbit IgG (H+L) at 1/10000 dilution

Performed under reducing conditions.

Predicted band size: 69 kDa

Observed band size: 75 kDa

Additional bands at: 110 kDa. We are unsure as to the identity of these extra bands.




Immunocytochemistry/ Immunofluorescence - Anti-PABP antibody (ab21060)

This image is courtesy of an anonymous Abreview

ab21060 at 1/50 staining HeLa (Human epithelial cell line from cervix adenocarcinoma) cells by ICC/IF.

HeLa cells fixed for 10 minutes with 1.9 % formaldehyde in 2 % sucrose in PBS then permeabilized for 10 minutes with 0.5% NP40 in PBS-10% sucrose. The cells were then blocked in horse serum and incubated with the antibody for 1 hour. An Alexa Fluor® 555 conjugated donkey anti-rabbit antibody was used as the secondary.




Immunocytochemistry/ Immunofluorescence - Anti-PABP antibody (ab21060)

This image is courtesy of an Abreview submitted by Sathya Thulasi Raman.

Immunocytochemical analysis of newborn porcine tracheal epithelial cells, labeling PABP with ab21060.

Cells were formaldehyde-fixed, permeabilized in cold methanol, and blocked with 5% horse serum in PBS for 40 minutes at 23°C.

Immunostaining with primary anti-PABP diluted 1/250 for 12 hours at 4°C.



Immunocytochemistry/ Immunofluorescence - Anti-PABP antibody (ab21060)

This image is courtesy of an anonymous Abreview.

Immunocytochemical analysis of rat OLN93 oligodendroglia cells, labeling PABP with ab21060.

Cells were paraformaldehyde-fixed, permeabilized with 0.1% Triton X-100, and blocked with 10% normal goat serum for 1 hour at 20°C.

Immunostaining with anti-PABP diluted 1/200 overnight at 4°C.

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors