

Anti-PADI2 / PAD2 antibody ab16478

★★★★★ [2 Abreviews](#) [14 References](#) [2 图像](#)

概述

产品名称	Anti-PADI2 / PAD2抗体
描述	兔多克隆抗体to PADI2 / PAD2
宿主	Rabbit
特异性	<p>ab16478 recognises a specific 75kDa band corresponding to PADI2, which is specifically blocked using the immunizing peptide in human colon, skeletal muscle and kidney lysates. There is a non-specific 18kDa band present in skeletal muscle lysates, which is attributed to cross-reactivity of the PADI2 antibody.</p> <p>Replenishment batches of ab16478 are tested in WB. Previous batches were additionally validated in ELISA, Flow Cyt, ICC/IF and IHC-P. These applications are still expected to work and are covered by our Abpromise guarantee.</p>
经测试应用	适用于: IHC-P, ELISA, ICC/IF, Flow Cyt, WB
种属反应性	与反应: Rat, Human
免疫原	<p>Synthetic peptide corresponding to Human PADI2/ PAD2 aa 100-200 (internal sequence) conjugated to keyhole limpet haemocyanin.</p> <p>(Peptide available as ab17091)</p>
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	<p>pH: 7.40</p> <p>Preservative: 0.02% Sodium azide</p> <p>Constituent: PBS</p>

Batches of this product that have a concentration < 1mg/ml may have BSA added as a stabilising agent. If you would like information about the formulation of a specific lot, please contact our scientific support team who will be happy to help.

纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab16478于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-P		Use at an assay dependent concentration.
ELISA		Use at an assay dependent concentration. PubMed: 19085382
ICC/IF		Use a concentration of 5 µg/ml.
Flow Cyt	★★★★★ (1)	Use at an assay dependent concentration. ab171870 - Rabbit polyclonal IgG, is suitable for use as an isotype control with this antibody.
WB	★★★☆☆ (1)	Use a concentration of 1 µg/ml. Detects a band of approximately 75 kDa (predicted molecular weight: 76 kDa).

靶标

功能	Catalyzes the deimination of arginine residues of proteins.
序列相似性	Belongs to the protein arginine deiminase family.
细胞定位	Cytoplasm.

图片



Anti-PADI2 / PAD2 antibody (ab16478) at 1 µg/ml + Human kidney tissue lysate - total protein ([ab30203](#)) at 20 µg

Secondary

Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/10000 dilution

Developed using the ECL technique.

Performed under reducing conditions.


Predicted band size: 76 kDa

Observed band size: 75 kDa

Additional bands at: 34 kDa. We are unsure as to the identity of these extra bands.

Exposure time: 20 minutes

This blot was produced using a 10% Bis-tris gel under the MOPS buffer system. The gel was run at 200V for 50 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour using 5% Bovine Serum Albumin before being incubated with ab16478 overnight at 4°C. Antibody binding was detected using an anti-rabbit antibody conjugated to HRP, and visualised using ECL development solution. The 75-kDa band observed is consistent with what has been described in the literature (PMID:18668562; 20668670; 16723463).



Immunocytochemistry/ Immunofluorescence - Anti-PADI2 / PAD2 antibody (ab16478)

ICC/IF image of ab16478 stained human Hek293 cells. The cells were PFA fixed (10 min), permeabilised in 0.1% PBS-Tween (20 min) and incubated with the antibody (ab16478, 5μg/ml) for 1h at room temperature. 1%BSA / 10% normal goat serum / 0.3M glycine was used to block non-specific protein-protein interactions. The secondary antibody (green) was Alexa Fluor® 488 goat anti-rabbit IgG (H+L) used at a 1/1000 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors