


Anti-PAPD5 antibody ab118341

★★★★★ [1 Abreviews](#) [1 图像](#)

概述

产品名称	Anti-PAPD5抗体
描述	山羊多克隆抗体to PAPD5
宿主	Goat
特异性	ab118341 is expected to recognize both reported isoforms (NP_001035374.2; NP_001035375.2).
经测试应用	适用于: WB
种属反应性	与反应: Mouse 预测可用于: Horse, Cow, Dog, Human 
免疫原	Synthetic peptide corresponding to Human PAPD5 aa 450-550 (internal sequence) (Cysteine residue). NP_001035374.2; NP_001035375.2 Run BLAST with ExPASy Run BLAST with NCBI
阳性对照	WB: Mouse and rat spleen lysate.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid repeated freeze / thaw cycles.
存储溶液	pH: 7.30 Preservative: 0.02% Sodium azide Constituents: 0.5% BSA, Tris buffered saline
纯度	Immunogen affinity purified
纯化说明	ab118341 is purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

克隆 多克隆
同种型 IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab118341于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。


应用	Ab评论	说明
WB	★☆☆☆☆ (1)	Use a concentration of 0.2 - 0.6 µg/ml. Detects a band of approximately 63 kDa (predicted molecular weight: 63 kDa). 1 hour primary incubation is recommended for this product.

靶标

相关性 Plays a role in replication-dependent histone mRNA degradation. May be involved in the terminal uridylation of mature histone mRNAs before their degradation is initiated. DNA polymerase, probably involved in DNA repair. May play a role in sister chromatid cohesion

细胞定位 Nuclear

图片



Anti-PAPD5 antibody (ab118341) at 0.2 µg/ml (for 1 hour) + Mouse spleen lysate (in RIPA buffer) at 35 µg

Developed using the ECL technique.

Predicted band size: 63 kDa

Observed band size: 70 kDa

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you

- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors