

Anti-PARK7/DJ1 antibody [EP2815Y] ab76008

敲除验证
重组
RabMAb

★★★★★
[3 Abreviews](#)
[38 References](#)
[13 图像](#)

概述

产品名称	Anti-PARK7/DJ1抗体[EP2815Y]
描述	兔单克隆抗体[EP2815Y] to PARK7/DJ1
宿主	Rabbit
特异性	The mouse and rat recommendation is based on the WB results. We do not guarantee IHC-P for mouse and rat.
经测试应用	适用于: WB, IP, IHC-P, Flow Cyt (Intra), ICC/IF
种属反应性	与反应: Mouse, Rat, Human
免疫原	Synthetic peptide corresponding to Human PARK7/DJ1 aa 1-100 (N terminal). Database link: Q99497
阳性对照	WB: Jurkat, HeLa, NIH3T3 or 293T cell lysate. Human fetal brain; Human brain nuclear fraction tissue lysate; Mouse brain and Rat brain tissue lysates. IHC-P: Human Lung and Brain tissue. ICC/IF: PANC-1 and Jurkat cell lines. Flow Cyt (intra): HepG2 cells. IP: Mouse brain lysate.
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
存储溶液	<p>pH: 7.20</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: 59% PBS, 40% Glycerol, 0.05% BSA</p>
纯度	Protein A purified

克隆单克隆

克隆编号EP2815Y

同种型IgG

应用

The Abpromise guarantee

Abpromise™承诺保证使用ab76008于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB	★★★★★ (2)	1/5000. Predicted molecular weight: 20 kDa. For unpurified use at 1/10000 - 1/20000.
IP		1/20.
IHC-P		1/1000. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. See IHC antigen retrieval protocols. Perform heat mediated antigen retrieval using 0.01M Sodium Citrate Buffer, pH 6.0 before commencing with IHC staining protocol.
Flow Cyt (Intra)		1/20. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody. For unpurified use at 1/100
ICC/IF	★★★★★ (1)	1/50 - 1/500.

靶标

功能

Protects cells against oxidative stress and cell death. Plays a role in regulating expression or stability of the mitochondrial uncoupling proteins SLC25A14 and SLC25A27 in dopaminergic neurons of the substantia nigra pars compacta and attenuates the oxidative stress induced by calcium entry into the neurons via L-type channels during pacemaking. Eliminates hydrogen peroxide and protects cells against hydrogen peroxide-induced cell death. May act as an atypical peroxiredoxin-like peroxidase that scavenges hydrogen peroxide. Following removal of a C-terminal peptide, displays protease activity and enhanced cytoprotective action against oxidative stress-induced apoptosis. Stabilizes NFE2L2 by preventing its association with KEAP1 and its subsequent ubiquitination. Binds to OTUD7B and inhibits its deubiquitinating activity. Enhances RELA nuclear translocation. Binds to a number of mRNAs containing multiple copies of GG or CC motifs and partially inhibits their translation but dissociates following oxidative stress. Required for correct mitochondrial morphology and function and for autophagy of dysfunctional mitochondria. Regulates astrocyte inflammatory responses. Acts as a positive regulator of androgen receptor-dependent transcription. Prevents aggregation of SNCA. Plays a role in fertilization. Has no proteolytic activity. Has cell-growth promoting activity and transforming activity. May function as a redox-sensitive chaperone.

组织特异性

Highly expressed in pancreas, kidney, skeletal muscle, liver, testis and heart. Detected at slightly

lower levels in placenta and brain. Detected in astrocytes, Sertoli cells, spermatogonia, spermatids and spermatozoa.

疾病相关

Defects in PARK7 are the cause of Parkinson disease type 7 (PARK7) [MIM:606324]. A neurodegenerative disorder characterized by resting tremor, postural tremor, bradykinesia, muscular rigidity, anxiety and psychotic episodes. PARK7 has onset before 40 years, slow progression and initial good response to levodopa. Some patients may show traits reminiscent of amyotrophic lateral sclerosis-parkinsonism/dementia complex (Guam disease).

序列相似性

Belongs to the peptidase C56 family.

翻译后修饰

Sumoylated on Lys-130 by PIAS2 or PIAS4; which is enhanced after ultraviolet irradiation and essential for cell-growth promoting activity and transforming activity.


Cys-106 is easily oxidized to sulfinic acid.

Undergoes cleavage of a C-terminal peptide and subsequent activation of protease activity in response to oxidative stress.

细胞定位

Cytoplasm. Nucleus. Mitochondrion. Under normal conditions, located predominantly in the cytoplasm and, to a lesser extent, in the nucleus and mitochondrion. Translocates to the mitochondrion and subsequently to the nucleus in response to oxidative stress and exerts an increased cytoprotective effect against oxidative damage. Detected in tau inclusions in brains from neurodegenerative disease patients.

图片



All lanes : Anti-PARK7/DJ1 antibody [EP2815Y] (ab76008) at 1/1000 dilution

Lane 1 : Wild-type HEK-293T (Human epithelial cell line from embryonic kidney transformed with large T antigen) whole cell lysate

Lane 2 : PARK7 knockout HEK-293T (Human epithelial cell line from embryonic kidney transformed with large T antigen) whole cell lysate

Lane 3 : HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate

Lane 4 : Human brain nuclear fraction tissue lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) at 1/10000 dilution


Predicted band size: 20 kDa

Observed band size: 24 kDa

Lanes 1-4: Merged signal (red and green). Green - ab76008


observed at 24 kDa. Red - loading control **ab8245** observed at 36 kDa.

ab76008 Anti-PARK7/DJ1 antibody [EP2815Y] was shown to specifically react with PARK7/DJ1 in wild-type HEK-293T cells. Loss of signal was observed when knockout cell line **ab266338** (knockout cell lysate **ab257016**) was used. Wild-type and PARK7/DJ1 knockout samples were subjected to SDS-PAGE. ab76008 and Anti-GAPDH antibody [6C5] - Loading Control (**ab8245**) were incubated overnight at 4°C at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.




Immunocytochemistry/ Immunofluorescence analysis of Jurkat (Human T cell leukemia T lymphocyte) cells labeling PARK7/DJ1 with Purified ab76008 at 1:500 dilution (0.2 µg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with Ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1:200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) was used as the secondary antibody at 1:1000 (2 µg/ml) dilution. DAPI nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.

Immunocytochemistry/ Immunofluorescence - Anti-PARK7/DJ1 antibody [EP2815Y] (ab76008)




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PARK7/DJ1 antibody [EP2815Y] (ab76008)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human lung cancer tissue sections labeling PARK7/DJ1 with Purified ab76008 at 1:1000 dilution (0.11 µg/ml). Heat mediated antigen retrieval was performed using [ab93684](#) (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Flow Cytometry (Intracellular) - Anti-PARK7/DJ1 antibody [EP2815Y] (ab76008)

Intracellular Flow Cytometry analysis of HepG2 (Human hepatocellular carcinoma epithelial cell) cells labeling PARK7/DJ1 with Purified ab76008 at 1/20 dilution (10 µg/ml) (red). Cells were fixed with 4% Paraformaldehyde. A Goat anti rabbit IgG (Alexa Fluor® 488, [ab150077](#)) secondary antibody was used at 1/2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).



Western blot - Anti-PARK7/DJ1 antibody [EP2815Y] (ab76008)

Lane 1: Wild-type HAP1 cell lysate (20 µg)
Lane 2: PARK7/DJ1 knockout HAP1 cell lysate (20 µg)
Lane 3: HeLa cell lysate (20 µg)
Lane 4: Human brain tissue lysate (20 µg)
Lanes 1 - 4: Merged signal (red and green). Green - ab76008 observed at 24 kDa. Red - loading control, [ab8245](#), observed at 37 kDa.


ab76008 was shown to specifically react with PARK/DJ1 in wild-type HAP1 cells. No band was observed when PARK/DJ1 knockout samples were used. Wild-type and PARK/DJ1 knockout samples were subjected to SDS-PAGE. ab76008 and [ab8245](#) (loading control to GAPDH) were both diluted 1/10,000 and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG

H&L (IRDye® 800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1/10,000 dilution for 1 h at room temperature before imaging.



Immunoprecipitation - Anti-PARK7/DJ1 antibody [EP2815Y] (ab76008)

ab76008 (purified) at 1:20 dilution (0.5µg) immunoprecipitating PARK7/DJ1 in Mouse brain lysate.
 Lane 1 (input): Mouse brain lysate 10µg
 Lane 2 (+): ab76008 & Mouse brain lysate
 Lane 3 (-): Rabbit monoclonal IgG (**ab172730**) instead of ab76008 in Mouse brain lysate
 For western blotting, VeriBlot for IP Detection Reagent (HRP) (**ab131366**) was used for detection at 1:1000 dilution.
 Blocking and diluting buffer: 5% NFDm/TBST.



Western blot - Anti-PARK7/DJ1 antibody [EP2815Y] (ab76008)

All lanes : Anti-PARK7/DJ1 antibody [EP2815Y] (ab76008) at 1/5000 dilution (Purified)

Lane 1 : HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates

Lane 2 : Mouse brain lysates

Lane 3 : Rat brain lysates


Lysates/proteins at 15 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/20000 dilution

Predicted band size: 20 kDa

Observed band size: 23 kDa



Western blot - Anti-PARK7/DJ1 antibody [EP2815Y] (ab76008)


Anti-PARK7/DJ1 antibody [EP2815Y] (ab76008) at 1/5000 dilution (Purified) + Human fetal brain lysates at 15 µg

Secondary

Goat Anti-Rabbit IgG (HRP) with minimal cross-reactivity with human IgG at 1/2000 dilution

Predicted band size: 20 kDa

Observed band size: 23 kDa




Immunocytochemistry/ Immunofluorescence - Anti-PARK7/DJ1 antibody [EP2815Y] (ab76008)

Immunocytochemistry/Immunofluorescence analysis of MCF-7 (human breast carcinoma) labelling PARK7/DJ1 with purified ab76008 at 1/500. Cells were fixed with 4% PFA and permeabilized with 0.1% Triton X-100. An Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/1000) was used as the secondary antibody (Ab150077). Nuclei counterstained with DAPI (blue).

Control: PBS only


This image was generated using the unpurified version of the product.



Flow Cytometry (Intracellular) - Anti-PARK7/DJ1 antibody [EP2815Y] (ab76008)

Overlay histogram showing HepG2 cells stained with ab76008 (red line). The cells were fixed with 4% paraformaldehyde (10 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab76008, 1/100 dilution) for 30 min at 22°C. The secondary antibody used was Alexa Fluor® 488 goat anti-rabbit IgG (H+L) ([ab150077](#)) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (1 µg/1x10⁶ cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter. This antibody gave a positive signal in HepG2 cells fixed with 80% methanol (5 min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.

This image was generated using the unpurified version of the product.



Western blot - Anti-PARK7/DJ1 antibody [EP2815Y] (ab76008)

All lanes : Anti-PARK7/DJ1 antibody [EP2815Y] (ab76008) at 1/20000 dilution

Lane 1 : Jurkat cell lysate

Lane 2 : HeLa cell lysate

Lane 3 : NIH3T3 cell lysate

Lane 4 : 293T cell lysate

Lysates/proteins at 10 µg per lane.


Secondary

All lanes : Goat anti-rabbit HRP at 1/1000 dilution

Predicted band size: 20 kDa

Observed band size: 23 kDa

This image was generated using the unpurified version of the product.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PARK7/DJ1 antibody [EP2815Y] (ab76008)

ab76008, at 1/250 dilution, staining PARK7/DJ1 in human brain by immunohistochemistry using paraffin-embedded tissue.

This image was generated using the unpurified version of the product.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-PARK7/DJ1 antibody [EP2815Y] (ab76008)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise,

please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors