abcam

Product datasheet

PE Anti-CCR7 antibody [4B12] ab95669

1 References 1 图像

概述

产品名称 PE Anti-CCR7抗体[4B12]

描述 PE大鼠单克隆抗体[4B12] to CCR7

宿主 Rat

偶联物 PE. Ex: 488nm, Em: 575nm

特异性 ab95669 stains Mouse CCR7-GFP fusion protein-transfected RBL cells

经测试应用 适用于: Flow Cyt

种属反应性 与反应: Mouse

免疫原 Tissue, cells or virus corresponding to CCR7. CCR7 Rat basophilic leukaemia (RBL)-2H3

transfectants.

阳性对照 C57BL/6, Balb/C and SJL/J splenocytes and thymocytes.

常规说明 Excitation: 488 nm Emission: 575 nm.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

性能

形式 Liquid

存放说明 Shipped at 4°C. Store at +4°C.

存储溶液 pH: 7.20

Preservative: 0.09% Sodium azide

Constituents: 0.87% Sodium chloride, PBS

纯**度** Protein G purified

骨髓瘤 x63-Ag8.653

1

同种型 lgG2a 轻链类型 kappa

应用

The Abpromise guarantee Abpromise™承诺保证使用ab95669于以下的经测试应用


"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt		Use 1-5µg for 10 ⁶ cells. ab134674 - Rat monoclonal lgG2a, is suitable for use as an isotype control with this antibody.

靶标

功能	Receptor for the MIP-3-beta chemokine. Probable mediator of EBV effects on B-lymphocytes or of normal lymphocyte functions.
组织 特异性	Expressed in various lymphoid tissues and activated B- and T-lymphocytes, strongly up-regulated in B-cells infected with Epstein-Barr virus and T-cells infected with herpesvirus 6 or 7.
序列相似性	Belongs to the G-protein coupled receptor 1 family.
细胞定位	Cell membrane.

图片



Staining of C57BL/6 splenocytes with Anti-Mouse CD4 FITC and 0.5 μ g of ab95669 with staining carried out at 4 °C (left) or 37 °C (right). Cells in the lymphocyte gate were used for analysis

 $\textbf{Please note:} \ \ \textbf{All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"}$

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors