abcam

Product datasheet

PE Anti-CD200R antibody [OX108] ab33366

1 References 1 图像

概述

产品名称 PE Anti-CD200R抗体[OX108]

描述 PE小鼠单克隆抗体[OX108] to CD200R

宿主 Mouse

偶联物 PE. Ex: 488nm, Em: 575nm

经测试应用适用于: Flow Cyt种属反应性与反应: Human

免疫原 Fusion protein corresponding to Human CD200R.

Database link: Q8TD46

常规说明

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

性能

形式 Liquid

存放说明 Shipped at 4°C. Store at +4°C.

存储溶液 pH: 7.40

Preservative: 0.09% Sodium azide

Constituents: PBS, 1% BSA

纯**度** Protein G purified

纯**化**说明 Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

克隆 单克隆

克隆编号 OX108

同种型 lgG1

田 紡

1

The Abpromise guarantee

Abpromise™承诺保证使用ab33366于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt		Use at an assay dependent concentration. Use neat - 1/10. Use 10ul working dilution to label 106 cells in 100ul


靶标

功能	Receptor for the CD200/OX2 cell surface glycoprotein. Also binds to HHV-8 K14 viral CD200 homolog with identical affinity and kinetics as the host CD200.
组织 特异性	Expressed in granulocytes, monocytes, most T cells, neutrophils, basophils and a subset of NK, NKT and B cells (at protein level). Expressed in bone marrow, lymph nodes, spleen, lung, liver, spinal cord, kidney. Expressed in monocyte-derived dendritic and mast cells.
序列相似性	Belongs to the CD200R family. Contains 1 lg-like C2-type (immunoglobulin-like) domain. Contains 1 lg-like V-type (immunoglobulin-like) domain.
翻译后修饰	The mature form of isoform 2 and/or isoform 4 starts at sequence position 27 of the corresponding isoform.

Secreted and Cell membrane.

图片

细胞定位



Staining of human peripheral blood lymphocytes with ab33366

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors