

PE Anti-CD86 antibody [BU63] ab77226

[4 References](#) [2 图像](#)

概述

产品名称	PE Anti-CD86抗体[BU63]
描述	PE小鼠单克隆抗体[BU63] to CD86
宿主	Mouse
偶联物	PE. Ex: 488nm, Em: 575nm
特异性	ab77226 reacts with CD86 expressed on professional antigen-presenting cells, such as dendritic cells, macrophages or activated B lymphocytes.
经测试应用	适用于: Flow Cyt
种属反应性	与反应: Human
免疫原	Tissue, cells or virus corresponding to Human CD86. B-lymphoblastoid cell line ARH 77
常规说明	<p>The antibody is conjugated with R-Phycoerythrin (PE) under optimum conditions.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C.
存储溶液	<p>pH: 7.4</p> <p>Preservative: 0.097% Sodium azide</p> <p>Constituents: 0.2% BSA, PBS</p>
纯度	Size exclusion
纯化说明	The conjugate is purified by size-exclusion chromatography and adjusted for direct use.
克隆	单克隆
克隆编号	BU63
同种型	IgG1

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab77226于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt		

应用说明

Flow Cyt: The reagent is designed for Flow Cytometry analysis of human blood cells
Use 20 µl/100 µl of whole blood or 10⁶ cells in a suspension.

Not yet tested in other applications.
Optimal dilutions/concentrations should be determined by the end user.

靶标

功能

Receptor involved in the costimulatory signal essential for T-lymphocyte proliferation and interleukin-2 production, by binding CD28 or CTLA-4. May play a critical role in the early events of T-cell activation and costimulation of naive T-cells, such as deciding between immunity and anergy that is made by T-cells within 24 hours after activation. Isoform 2 interferes with the formation of CD86 clusters, and thus acts as a negative regulator of T-cell activation.

组织特异性

Expressed by activated B-lymphocytes and monocytes.

序列相似性

Contains 1 Ig-like C2-type (immunoglobulin-like) domain.
Contains 1 Ig-like V-type (immunoglobulin-like) domain.


翻译后修饰

Polyubiquitinated; which is promoted by MARCH8 and results in endocytosis and lysosomal degradation.

细胞定位


Membrane.

图片



Flow Cytometry - PE Anti-CD86 antibody [BU63]
(ab77226)

Analysis by flow cytometry of ab77226 staining CD86 in human peripheral blood cells.



Flow Cytometry - PE Anti-CD86 antibody [BU63]
(ab77226)

Flow cytometry analysis of human peripheral blood leukocytes with anti-human CD86 (BU63) PE.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors