abcam

Product datasheet

PE Anti-HDAC2 antibody [Y461] ab210827



重组 RabMAb

3 图像

概述

产品名称 PE Anti-HDAC2抗体[Y461]

描述 PE兔单克隆抗体[Y461] to HDAC2

宿主 Rabbit

偶联物 PE. Ex: 488nm, Em: 575nm

经测试应用 适用于: ICC/IF, Flow Cyt (Intra)

种属反应性 与反应: Human

预测可用于: Mouse, Rat 🔷

免疫原 Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

阳性对照 Flow Cyt (intra): HeLa cells ICC/IF: HeLa cells

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit 常规说明

monoclonal antibodies. For details on our patents, please refer to **RabMAb® patents**.

性能

形式 Liquid

存放说明 Shipped at 4°C. Store at 4°C (stable for up to 12 months). Upon delivery aliquot. Store at +4°C.

Do Not Freeze. Store In the Dark.

存储溶液 pH: 7.4

> Preservative: 0.02% Sodium azide Constituents: PBS, 1% BSA

纯度 Protein A purified

克隆 单克隆 克隆编号 Y461 同种型 ΙgG

应用

"应用说明"部分 下显示的仅为推荐的起始稀释度:实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC/IF		1/100. This product gave a positive signal in HeLa cells fixed with 4% formaldehyde (10 min) and 100% methanol (5 min)
Flow Cyt (Intra)		1/500.

靶标

功能

Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes.

Forms transcriptional repressor complexes by associating with MAD, SIN3, YY1 and N-COR. Interacts in the late S-phase of DNA-replication with DNMT1 in the other transcriptional repressor complex composed of DNMT1, DMAP1, PCNA, CAF1. Deacetylates TSHZ3 and regulates its transcriptional repressor activity.

组织**特异性** Widely expressed; lower levels in brain and lung.


序列相似性 Belongs to the histone deacetylase family. HD type 1 subfamily.

翻译后修饰 S-nitrosylated by GAPDH. In neurons, S-Nitrosylation at Cys-262 and Cys-274 does not affect the enzyme activity but abolishes chromatin-binding, leading to increases acetylation of histones and activate genes that are associated with neuronal development. In embryonic cortical neurons, S-

Nitrosylation regulates dendritic growth and branching.

细胞定位 Nucleus.

图片




Flow Cytometry (Intracellular) - PE Anti-HDAC2 antibody [Y461] (ab210827)

Overlay histogram showing HeLa cells stained with ab210827 (red line). The cells were fixed with 4% formaldehyde (10 min) and then permeabilized with 90% methanol for 30 min at -20°C. The cells were then incubated in 1x PBS / 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody (ab210827, 1/500 dilution) for 30 min at 22°C.

Isotype control antibody (black line) was rabbit IgG (monoclonal) Phycoerythrin (ab209478) used at the same concentration and conditions as the primary antibody. Unlabelled sample (blue line) was also used as a control.

Acquisition of >5,000 events were collected using a 50mW Yellow-Green laser (561nm) and 586/15 bandpass filter.




Immunocytochemistry/ Immunofluorescence - PE
Anti-HDAC2 antibody [Y461] (ab210827)

ab210827 staining HDAC2 in HeLa cells. The cells were fixed with 4% formaldehyde (10 min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab210827 at 1/100 dilution (pseudocolored in green) and ab195884, Rat monoclonal to Tubulin (Alexa Fluor® 647), at 1/250 dilution (shown in red). Nuclear DNA was labelled with DAPI (shown in blue).

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

This product also gave a positive signal under the same testing conditions in HeLa cells fixed with 100% methanol (5min).



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors