# abcam

**Product datasheet** 

## PE Anti-Prohibitin antibody [EP2803Y] ab210424

重组 RabMAb

### <u>1 Abreviews</u> 3 图像

概述			
产品名称	PE Anti-Prohibitin <b>抗体</b> [EP2803Y]		
描述	PE兔单克隆抗体[EP2803Y] to Prohibitin		
宿主	Rabbit		
偶联物	PE. Ex: 488nm, Em: 575nm		
经测试应 <b>用</b>	适用于: Flow Cyt (Intra), ICC/IF		
种属反应性	与反应: Human		
	预测可用于: Mouse, Rat 🛛 📤		
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.		
<b>阳性</b> 对照	Flow Cyt (intra): HeLa cells ICC/IF: HeLa cells		
<b>常</b> 规说 <b>明</b>	<ul> <li>This product is a recombinant monoclonal antibody, which offers several advantages including:</li> <li>High batch-to-batch consistency and reproducibility</li> <li>Improved sensitivity and specificity</li> <li>Long-term security of supply</li> <li>Animal-free production</li> <li>For more information <u>see here</u>.</li> <li>Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <u>RabMAb<sup>®</sup> patents</u>.</li> </ul>		
性能			
形式	Liquid		
存 <b>放</b> 说明	Shipped at 4°C. Upon delivery aliquot. Store at +4°C. Do Not Freeze. Store In the Dark.		
存储溶液	pH: 7.4 Preservative: 0.02% Sodium azide		

纯度 Protein A purified

lgG

Constituents: PBS, 1% BSA

**克隆** 单克隆

**克隆编号** EP2803Y

同种型

#### 应用


#### The Abpromise guarantee Abpromise ™承诺保证使用ab210424于以下的经测试应用

#### "应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt (Intra)		1/500.
ICC/IF		1/100. This product gave a positive signal in HeLa cells fixed with 4% formaldehyde (10 min) and 100% methanol (5 min)

<b>靶</b> 标	
功能	Prohibitin inhibits DNA synthesis. It has a role in regulating proliferation. As yet it is unclear if the protein or the mRNA exhibits this effect. May play a role in regulating mitochondrial respiration activity and in aging.
组织 <b>特异性</b>	Widely expressed in different tissues.
序列相似性	Belongs to the prohibitin family.
发 <b>展</b> 阶 <b>段</b>	Levels of expression in fibroblasts decrease heterogeneously during cellular aging.
细 <b>胞定位</b>	Mitochondrion inner membrane.


#### 图片



Flow Cytometry (Intracellular) - PE Anti-Prohibitin antibody [EP2803Y] (ab210424) Overlay histogram showing HeLa cells stained with ab210424 (red line). The cells were fixed with 4% formaldehyde (10 min) and then permeabilized with 90% methanol for 30 min at -20°C. The cells were then incubated in 1x PBS / 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody (ab210424, 1/500 dilution) for 30 min at 22°C.

Isotype control antibody (black line) was rabbit IgG (monoclonal) Phycoerythrin (<u>ab209478</u>) used at the same concentration and conditions as the primary antibody. Unlabelled sample (blue line) was also used as a control.

Acquisition of >5,000 events were collected using a 50mW Yellow-Green laser (561nm) and 586/15 bandpass filter.



Immunocytochemistry/ Immunofluorescence - PE Anti-Prohibitin antibody [EP2803Y] (ab210424)



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise,

ab210424 staining Prohibitin in HeLa cells. The cells were fixed with 4% formaldehyde (10 min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab210424 at 1/100 dilution (Pseudocolored in green) and **ab195884**, Rat monoclonal to Tubulin (Alexa Fluor<sup>®</sup> 647), at 1/250 dilution (shown in red). Nuclear DNA was labelled with DAPI (shown in blue).

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

This product also gave a positive signal under the same testing conditions in HeLa cells fixed with 100% methanol (5min).

please visit https://www.abcam.cn/abpromise or contact our technical team.

#### **Terms and conditions**

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors