

### Anti-Perforin antibody [CB5.4] ab16074

★★★★★ **5 Abreviews** **29 References**

#### 概述

<b>产品名称</b>	Anti-Perforin抗体[CB5.4]
<b>描述</b>	大鼠单克隆抗体[CB5.4] to Perforin
<b>宿主</b>	Rat
<b>经测试应用</b>	<b>适用于:</b> IP, ICC, WB, IHC-Fr, IHC-P, ICC/IF
<b>种属反应性</b>	<b>与反应:</b> Mouse
<b>免疫原</b>	Recombinant fragment, corresponding to amino acids 98 - 534 of Mouse Perforin.
<b>表位</b>	This antibody recognizes mouse perforin, region amino acids 402-534.
<b>常规说明</b>	<p>This product was changed from ascites to tissue culture supernatant on 31<sup>st</sup> May 2019. Please note that the dilutions may need to be adjusted accordingly. If you have any questions, please do not hesitate to contact our scientific support team.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 性能

<b>形式</b>	Liquid
<b>存放说明</b>	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid repeated freeze / thaw cycles.
<b>存储溶液</b>	Preservative: 0.02% Sodium azide Constituent: PBS
<b>纯度</b>	Tissue culture supernatant
<b>克隆</b>	单克隆
<b>克隆编号</b>	CB5.4
<b>同种型</b>	IgG2a

**The Abpromise guarantee**      **Abpromise™**承诺保证使用ab16074于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IP		Use at an assay dependent concentration.
ICC		Use at an assay dependent concentration.
WB	★★★★★ (1)	Use at an assay dependent concentration.
IHC-Fr		Use at an assay dependent concentration.
IHC-P	★★★★☆ (3)	Use at an assay dependent concentration.
ICC/IF	★★★★☆ (1)	Use at an assay dependent concentration.

**靶标**

<b>功能</b>	Plays a key role in secretory granule-dependent cell death, and in defense against virus-infected or neoplastic cells. Plays an important role in killing other cells that are recognized as non-self by the immune system, e.g. in transplant rejection or some forms of autoimmune disease. Can insert into the membrane of target cells in its calcium-bound form, oligomerize and form large pores. Promotes cytolysis and apoptosis of target cells by facilitating the uptake of cytotoxic granzymes.
<b>疾病相关</b>	Defects in PRF1 are the cause of hemophagocytic lymphohistiocytosis familial type 2 (FHL2) [MIM:603553]; also known as HPLH2. Familial hemophagocytic lymphohistiocytosis (FHL) is a genetically heterogeneous, rare autosomal recessive disorder. It is characterized by immune dysregulation with hypercytokinemia and defective natural killer cell function. The clinical features of the disease include fever, hepatosplenomegaly, cytopenia, hypertriglyceridemia, hypofibrinogenemia, and neurological abnormalities ranging from irritability and hypotonia to seizures, cranial nerve deficits, and ataxia. Hemophagocytosis is a prominent feature of the disease, and a non-malignant infiltration of macrophages and activated T lymphocytes in lymph nodes, spleen, and other organs is also found.
<b>序列相似性</b>	Belongs to the complement C6/C7/C8/C9 family. Contains 1 C2 domain. Contains 1 EGF-like domain. Contains 1 MACPF domain.
<b>结构域</b>	The C2 domain mediates calcium-dependent binding to lipid membranes. A subsequent conformation change leads to membrane insertion of beta-hairpin structures and pore formation. The pore is formed by transmembrane beta-strands.
<b>翻译后修饰</b>	N-glycosylated.
<b>细胞定位</b>	Cytoplasmic granule lumen. Secreted. Cell membrane. Endosome lumen. Stored in cytoplasmic granules of cytolytic T-lymphocytes and secreted into the cleft between T-lymphocyte and target cell. Inserts into the cell membrane of target cells and forms pores. Membrane insertion and pore formation requires a major conformation change. May be taken up via endocytosis involving clathrin-coated vesicles and accumulate in a first time in large early endosomes.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors