# abcam

### Product datasheet

## Anti-PIAS1 + PIAS2 antibody [EPR2581Y] ab77231



重组 RabMAb

★★★★★ 4 Abreviews 18 References 5 图像

概述

产品名称 Anti-PIAS1 + PIAS2抗体[EPR2581Y]

描述 兔单克隆抗体[EPR2581Y] to PIAS1 + PIAS2

宿主 Rabbit

经测试应用 适用于: ICC/IF, WB, IHC-P

种属反应性 与反应: Human, Recombinant fragment

预测可用于: Mouse, Rat 🔷

免疫原 Synthetic peptide within Human PIAS1 aa 600 to the C-terminus (C terminal). The exact

> sequence is proprietary. Database link: **O75925**

阳性对照 Daudi, 293, Jurkat, SW40 and HepG2 cell lysates; human tonsil tissue.

常规说明 This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity - Long-term security of supply

- Animal-free production

For more information **see here**.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

性能

形式

存放说明 Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

存储溶液 pH: 7.20

Preservative: 0.01% Sodium azide

Constituents: 9% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA, 50% Tissue culture

supernatant

纯度 Protein A purified

克隆 单克隆

#### 应用

#### The Abpromise guarantee Abpromise™承诺保证使用ab77231于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。


应用	Ab评论	说明
ICC/IF	<b>★★★★</b> <u>(1)</u>	1/100 - 1/500.
WB	**** (1)	1/1000 - 1/10000. Predicted molecular weight: 72 kDa.
IHC-P		1/100 - 1/250. Perform heat mediated antigen retrieval via the pressure cooker method before commencing with IHC staining protocol.

#### 靶标

#### 细胞定位

PIAS1: Nucleus speckle. Interaction with CSRP2 may induce a partial redistribution along the cytoskeleton. PIAS2: Nucleus speckle. Nucleus, PML body. Nucleus. Colocalizes at least partially with promyelocytic leukemia nuclear bodies (PML NBs) (PubMed:22406621). Colocalizes with SUMO1 in nuclear granules (By similarity).

#### 图片



**All lanes :** Anti-PIAS1 + PIAS2 antibody [EPR2581Y] (ab77231) at 1/50000 dilution

**Lane 1**: Human PIAS1 full-length recombinant protein with GST-Tag

**Lane 2**: Human PIAS2-beta recombinant protein fragment with GST-His-Tag

**Lane 3 :** Human PIAS3 recombinant protein fragment with SUMO-His-Tag

Lysates/proteins at 0.015 µg per lane.

#### Secondary

**All lanes :** Goat Anti-Rabbit lgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 72 kDa

Observed band size: 98, 42 kDa

Blocking and Diluting buffer and concentration: 5% NFDM /TBST

Exposure time: Lane1:10 seconds, Lane2: 5 seconds, Lane3: 3 minutes.


Human PIAS1 full-length recombinant protein (Cat#: ab152888)

**All lanes**: Anti-PIAS1 + PIAS2 antibody [EPR2581Y] (ab77231) at 1/50000 dilution

Lane 1 : Daudi cell lysate
Lane 2 : 293 cell lysate
Lane 3 : Jurkat cell lysate
Lane 4 : SW40 cell lysate

Lane 5: HepG2 cell lysate

Lysates/proteins at 10 µg per lane.



[EPR2581Y] (ab77231)

5

kDa

150-

100-

75-

50-


37-

25-

## Secondary

All lanes: HRP labelled goat anti-rabbit at 1/2000 dilution


**Predicted band size:** 72 kDa **Observed band size:** 78 kDa



Immunocytochemistry/ Immunofluorescence - Anti-PIAS1 + PIAS2 antibody [EPR2581Y] (ab77231)

Immunocytochemistry/Immunofluorescence analysis of HeLa (human cervix adenocarcinoma) cells labelling PIAS1 + PIAS2 with purified ab77231 at 1/500. Cells were fixed with 4% Paraformaldehyde and permeabilized with 0.1% Triton X-100. ab150077, Alexa Fluor<sup>®</sup> 488-conjugated goat anti-rabbit lgG (1/1000) was used as the secondary antibody. Nuclei were counterstained with DAPI (blue).


Secondary Only Control: PBS was used instead of the primary antibody as the negative control.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-PIAS1 + PIAS2 antibody [EPR2581Y] (ab77231)

ab77231, at a 1/100 dilution, staining PIAS1 + PIAS2 in paraffin embedded human tonsil tissue by Immunohistochemistry.

Perform heat mediated antigen retrieval via the pressure cooker method before commencing with IHC staining protocol.



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.cn/abpromise">https://www.abcam.cn/abpromise</a> or contact our technical team.

## Terms and conditions

Guarantee only valid for products bought direct from Abcam or one of our authorized distributors