

Anti-PP1C gamma antibody ab16387

5 References

概述

产品名称	Anti-PP1C gamma抗体
描述	羊多克隆抗体to PP1C gamma
宿主	Sheep
特异性	No cross reactivity with other recombinant pp1 isoforms.
经测试应用	适用于: IP, WB
种属反应性	与反应: Human 预测可用于: Mouse, Rat, Cow, Xenopus laevis 
免疫原	Synthetic peptide: TPPRGMITKQAKK conjugated to KLH, corresponding to amino acids 311-323 of Human PPP1G1.  Run BLAST with  Run BLAST with
阳性对照	Recombinant Human PP1C gamma protein (ab114828) can be used as a positive control in WB.
常规说明	<p>A protein phosphatase is a phosphatase enzyme that removes a phosphate group from the phosphorylated amino acid residue of its substrate protein. Protein phosphorylation is one of the most common forms of reversible protein posttranslational modification (PTM), with up to 30% of all proteins being phosphorylated at any given time. Protein kinases (PKs) are the effectors of phosphorylation and catalyse the transfer of a γ-phosphate from ATP to specific amino acids on proteins. Several hundred PKs exist in mammals and are classified into distinct super-families. Proteins are phosphorylated predominantly on Ser, Thr and Tyr residues, which account for 79.3, 16.9 and 3.8% respectively of the phosphoproteome, at least in mammals. In contrast, protein phosphatases (PPs) are the primary effectors of dephosphorylation and can be grouped into three main classes based on sequence, structure and catalytic function. The largest class of PPs is the phosphoprotein phosphatase (PPP) family comprising PP1, PP2A, PP2B, PP4, PP5, PP6 and PP7, and the protein phosphatase Mg^{2+} or Mn^{2+}-dependent (PPM) family, composed primarily of PP2C.</p> <p>Source: The immunogen used to generate the purified antibody was a peptide conjugated to KLH corresponding to the sequence NH₂-Thr-Pro-Pro-Arg-Gly-Met-Ile-Thr-Lys-Gln-Ala-Lys-Lys-COOH. This peptide antibody corresponds to C-terminal peptide of PP1 gamma 1 catalytic subunit having a MW of 37kD. The sequence used is amino acid 311-323.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p>

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
存储溶液	Preservative: 0.08% Sodium azide Constituent: PBS
纯度	Ammonium Sulphate Precipitation
克隆	多克隆
同种型	IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab16387于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IP		Use at an assay dependent concentration.
WB		Use at an assay dependent concentration.

靶标

功能	Protein phosphatase 1 (PP1) is essential for cell division, and participates in the regulation of glycogen metabolism, muscle contractility and protein synthesis. Involved in regulation of ionic conductances and long-term synaptic plasticity. May play an important role in dephosphorylating substrates such as the postsynaptic density-associated Ca(2+)/calmodulin dependent protein kinase II. Component of the PTW/PP1 phosphatase complex, which plays a role in the control of chromatin structure and cell cycle progression during the transition from mitosis into interphase.
序列相似性	Belongs to the PPP phosphatase family. PP-1 subfamily.
细胞定位	Cytoplasm. Nucleus. Nucleus > nucleolus. Nucleus > nucleoplasm. Nucleus speckle. Chromosome > centromere > kinetochore. Cleavage furrow. Midbody. Colocalizes with SPZ1 in the nucleus (By similarity). Rapidly exchanges between the nucleolar, nucleoplasmic and cytoplasmic compartments. Highly mobile in cells and can be relocalized through interaction with targeting subunits. In the presence of PPP1R8 relocalizes from the nucleolus to nuclear speckles. Shows a dynamic targeting to specific sites throughout the cell cycle. Highly concentrated in nucleoli of interphase cells and localizes at kinetochores early in mitosis. Relocalization to chromosome-containing regions occurs at the transition from early to late anaphase. Also accumulates at the cleavage furrow and midbody by telophase.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors