


# Anti-Proteasome 20S beta 6 antibody ab3331

**1 References**   **7 图像**

## 概述

产品名称	Anti-Proteasome 20S beta 6抗体
描述	兔多克隆抗体to Proteasome 20S beta 6
宿主	Rabbit
特异性	Detects proteasome 20S beta 6 from purified bovine and human 26S proteasome samples.
经测试应用	适用于: ICC/IF, WB
种属反应性	与反应: Mouse, Rat, Cow, Human 预测可用于: Xenopus laevis, Zebrafish 
免疫原	Synthetic peptide corresponding to Human Proteasome 20S beta 6 aa 41-58. Sequence: CRSGSAADTQAV/IADAVTY  Database link: <b>P28072</b> (Peptide available as <b>ab4947</b> )

 **Run BLAST with**

 **Run BLAST with**

阳性对照	WB: HeLa, BAEC whole cell lysates, HeLa, 3T3-L1, PC-3, MCF7, A549, PANC-1, Mouse Kidney, Mouse Liver and Rat Liver nuclear enriched cell lysate. ICC/IF: A431, HeLa, NIH 3T3, BAEC cells.
------	---

常规说明	<p>Abcam has not and does not intend to apply for the REACH Authorisation of customers' uses of products that contain European Authorisation list (Annex XIV) substances.</p> <p>It is the responsibility of our customers to check the necessity of application of REACH Authorisation, and any other relevant authorisations, for their intended uses.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>
------	--

## 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	Constituents: 3% Sodium deoxycholate, 3% Triton-X-100, 0.3% Tris HCl, 15% Sodium chloride
纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

应用

The Abpromise guarantee      **Abpromise™**承诺保证使用ab3331于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC/IF		1/10 - 1/100.
WB		Use a concentration of 1 µg/ml. Detects a band of approximately 25 kDa (predicted molecular weight: 25 kDa).

靶标

功能	The proteasome is a multicatalytic proteinase complex which is characterized by its ability to cleave peptides with Arg, Phe, Tyr, Leu, and Glu adjacent to the leaving group at neutral or slightly basic pH. The proteasome has an ATP-dependent proteolytic activity. This unit is responsible of the peptidyl glutamyl-like activity. May catalyze basal processing of intracellular antigens.
序列相似性	Belongs to the peptidase T1B family.
细胞定位	Cytoplasm. Nucleus.

图片



**All lanes :** Anti-Proteasome 20S beta 6 antibody (ab3331) at 1 µg/ml

**Lane 1 :** Nuclear enriched extracts from untransfected PC-3 cells


**Lane 2 :** Nuclear enriched extracts from non-targeting scrambled siRNA transfected PC-3 cells

**Lane 3 :** Nuclear enriched extracts from PSMB6 knockdown PC-3 cells

### Secondary

**All lanes :** Goat anti-Rabbit IgG (H+L) Superclonal™ Recombinant Secondary Antibody, HRP at 1/6000 dilution

**Predicted band size:** 25 kDa



**All lanes :** Anti-Proteasome 20S beta 6 antibody (ab3331) at 1 µg/ml

**Lane 1 :** HeLa nuclear enriched extract lysate

**Lane 2 :** 3T3-L1 nuclear enriched extract lysate

**Lane 3 :** PC-3 nuclear enriched extract lysate

**Lane 4 :** MCF7 nuclear enriched extract lysate

**Lane 5 :** A549 nuclear enriched extract lysate

**Lane 6 :** PANC-1 nuclear enriched extract lysate

**Lane 7 :** Mouse Kidney nuclear enriched extract lysate

**Lane 8 :** Mouse Liver nuclear enriched extract lysate

**Lane 9 :** Rat Liver nuclear enriched extract lysate

Lysates/proteins at 30 µg per lane.

### Secondary

**All lanes :** Goat anti-Rabbit IgG (H+L) Superclonal™ Recombinant Secondary Antibody, HRP at 1/6000 dilution


Developed using the ECL technique.

**Predicted band size:** 25 kDa

**Observed band size:** 21 kDa

Samples were electrophoresed using NuPAGE™ 12% Bis-Tris Protein Gel and the resolved proteins were then transferred onto a

Nitrocellulose membrane by iBlot® 2 Dry Blotting System. The blot was probed with the primary antibody (ab3331) and detected by chemiluminescence with secondary antibody using the iBright FL 1000. Chemiluminescent detection was performed using Novex® ECL Chemiluminescent Substrate Reagent Kit.



**All lanes** : Anti-Proteasome 20S beta 6 antibody (ab3331) at 1/1000 dilution

**Lane 1** : HeLa (Human epithelial adenocarcinoma cell line) whole cell lysate


**Lane 2** : BAEC (Bovine aortic endothelial cell line) whole cell lysate

Lysates/proteins at 25 µg per lane.


### Secondary

**All lanes** : HRP-conjugated secondary antibody

**Predicted band size:** 25 kDa




Immunocytochemistry/Immunofluorescence analysis of Proteasome 20S beta 6 (green) showing staining in the cytoplasm and nucleus of A431 (Human epidermoid carcinoma cell line) cells (right) compared to a negative control without primary antibody (left). Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with ab3331 in 3% BSA-PBS at a dilution of 1:20 and incubated overnight at 4°C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight-conjugated secondary antibody in PBS at room temperature in the dark. F-actin (red) was stained with a fluorescent red phalloidin and nuclei (blue) were stained with Hoechst or DAPI. Images were taken at a magnification of 60x.




Immunocytochemistry/ Immunofluorescence - Anti-Proteasome 20S beta 6 antibody (ab3331)

Immunofluorescent analysis of Proteasome 20S Y (green) in NIH/3T3 (Mouse embryo fibroblast cell line) cells. The cells were fixed with 4% paraformaldehyde for 15 minutes, permeabilized with 0.1% Triton X-100 in PBS for 15 minutes and blocked with 3% Blocker BSA in PBS for 30 minutes at room temperature. Cells were stained with or without Proteasome 20S Y rabbit polyclonal antibody, at a concentration of 5 µg/mL for 1 hour at room temperature, and then incubated with a Goat anti-Rabbit (H+L) Superclonal Secondary Antibody, Alexa Fluor® 488 conjugate at 1/1000 dilution for 1 hour at room temperature (both panels, green). Nuclei (both panels, blue) were stained with Hoechst 33342 dye. Images were taken at 20X magnification.



Immunocytochemistry/ Immunofluorescence - Anti-Proteasome 20S beta 6 antibody (ab3331)

Immunocytochemistry/Immunofluorescence analysis of Proteasome 20S beta 6 (green) showing staining in the cytoplasm and nucleus of BAEC (Bovine aortic endothelial cell line) cells (right) compared to a negative control without primary antibody (left). Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with ab3331 in 3% BSA-PBS at a dilution of 1:20 and incubated overnight at 4°C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight-conjugated secondary antibody in PBS at room temperature in the dark. F-actin (red) was stained with a fluorescent red phalloidin and nuclei (blue) were stained with Hoechst or DAPI. Images were taken at a magnification of 60x.



Immunocytochemistry/ Immunofluorescence - Anti-Proteasome 20S beta 6 antibody (ab3331)

Immunocytochemistry/Immunofluorescence analysis of Proteasome 20S beta 6 (green) showing staining in the cytoplasm and nucleus of HeLa (Human epithelial adenocarcinoma cell line) cells (right) compared to a negative control without primary antibody (left). Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with ab3331 in 3% BSA-PBS at a dilution of 1:20 and incubated overnight at 4°C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight-conjugated secondary antibody in PBS at room temperature in the dark. F-actin (red) was stained with a fluorescent red phalloidin and nuclei (blue) were stained with Hoechst or DAPI. Images were taken at a magnification of 60x.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors