


Anti-RAB7 antibody [Rab7-117] - Late Endosome Marker ab50533

★★★★☆ [27 Abreviews](#) [107 References](#) [7 图像](#)

概述

产品名称	Anti-RAB7抗体[Rab7-117] - Late Endosome Marker
描述	小鼠单克隆抗体[Rab7-117] to RAB7 - Late Endosome Marker
宿主	Mouse
经测试应用	适用于: ICC, WB
种属反应性	与反应: Mouse, Rat, Dog, Human 预测可用于: Rabbit, Chicken, Cow, Xenopus laevis, Monkey 
免疫原	Synthetic peptide: EQAQFTIARNALKQE , corresponding to amino acids 163-177 of Human RAB7 Run BLAST with Run BLAST with
阳性对照	WB: HeLA, HepG2, NRK , NIH3/T3 and MDCK cell lysates. ICC: HeLa cells.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	pH: 7.40 Preservative: 0.097% Sodium azide Constituent: 0.0268% PBS
纯度	IgG fraction
克隆	单克隆

克隆编号	Rab7-117
骨髓瘤	NS1
同种型	IgG2b

应用

The Abpromise guarantee **Abpromise™** 承诺保证使用ab50533于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC	★☆☆☆☆ (1)	Use at an assay dependent concentration.
WB	★★★★★ (18)	Use a concentration of 0.5 - 1 µg/ml. Predicted molecular weight: 23 kDa.

靶标

功能

Key regulator in endo-lysosomal trafficking. Governs early-to-late endosomal maturation, microtubule minus-end as well as plus-end directed endosomal migration and positioning, and endosome-lysosome transport through different protein-protein interaction cascades. Plays a central role, not only in endosomal traffic, but also in many other cellular and physiological events, such as growth-factor-mediated cell signaling, nutrient-transporter mediated nutrient uptake, neurotrophin transport in the axons of neurons and lipid metabolism. Also involved in regulation of some specialized endosomal membrane trafficking, such as maturation of melanosomes, pathogen-induced phagosomes (or vacuoles) and autophagosomes. Plays a role in the maturation and acidification of phagosomes that engulf pathogens, such as *S.aureus* and *M.tuberculosis*. Plays a role in the fusion of phagosomes with lysosomes. Plays important roles in microbial pathogen infection and survival, as well as in participating in the life cycle of viruses. Microbial pathogens possess survival strategies governed by RAB7A, sometimes by employing RAB7A function (e.g. *Salmonella*) and sometimes by excluding RAB7A function (e.g. *Mycobacterium*). In concert with RAC1, plays a role in regulating the formation of RBs (ruffled borders) in osteoclasts. Controls the endosomal trafficking and neurite outgrowth signaling of NTRK1/TRKA. Regulates the endocytic trafficking of the EGF-EGFR complex by regulating its lysosomal degradation.

组织特异性

Widely expressed; high expression found in skeletal muscle.

疾病相关

Defects in RAB7A are the cause of Charcot-Marie-Tooth disease type 2B (CMT2B) [MIM:600882]; also known as hereditary motor and sensory neuropathy II (HMSN2). CMT2B is a form of Charcot-Marie-Tooth disease, the most common inherited disorder of the peripheral nervous system. Charcot-Marie-Tooth disease is classified in two main groups on the basis of electrophysiologic properties and histopathology: primary peripheral demyelinating neuropathy or CMT1, and primary peripheral axonal neuropathy or CMT2. Neuropathies of the CMT2 group are characterized by signs of axonal regeneration in the absence of obvious myelin alterations, normal or slightly reduced nerve conduction velocities, and progressive distal muscle weakness and atrophy. CMT2B is clinically characterized by marked distal muscle weakness and a high frequency of foot ulcers, infections and amputations of the toes. CMT2B inheritance is autosomal dominant.


序列相似性

Belongs to the small GTPase superfamily. Rab family.

细胞定位

Late endosome. Lysosome. Cytoplasmic vesicle > phagosome. Melanosome. Cytoplasmic vesicle > phagosome membrane. Co-localizes with OSBPL1A at the late endosome. Found in the ruffled border (a late endosomal-like compartment in the plasma membrane) of bone-resorbing osteoclasts. Recruited to phagosomes containing *S.aureus* or *Mycobacterium*.

图片



All lanes : Anti-RAB7 antibody [Rab7-117] - Late Endosome Marker (ab50533) at 1 $\mu\text{g/ml}$

Lane 1 : HeLa cell lysate

Lane 2 : HepG2 cell lysate

Lane 3 : NRK (rat cell kidney line) cell lysate


Lane 4 : NIH/3T3 (mouse embryonic fibroblast cell line) cell lysate

Lane 5 : MDCK (dog kidney cell line) cell lysate


Secondary

All lanes : Goat Anti-Mouse IgG-peroxidase

Predicted band size: 23 kDa



Immunocytochemistry/ Immunofluorescence analysis of HeLa cells labeling RAB7 with ab50533 at 10 $\mu\text{g/mL}$. Cells were fixed and permeabilized with 4% paraformaldehyde followed by 0.5% saponin. Goat Anti-Mouse IgG, Atto-488 conjugate was used as the secondary antibody.



Western blot - Anti-RAB7 antibody [Rab7-117] - Late Endosome Marker (ab50533)

This image is courtesy of an Abreview submitted by Armen Petrosyan

Anti-RAB7 antibody [Rab7-117] - Late Endosome Marker (ab50533) at 1/2000 dilution + HepG2 whole cell lysate at 1/2000 dilution

Secondary

HRP-conjugated donkey anti-mouse IgG polyclonal at 1/1 dilution

Developed using the ECL technique.


Performed under reducing conditions.

Predicted band size: 23 kDa

Exposure time: 30 seconds

Blocked with 5% milk for 1 hour at 22°C.

Incubated with the primary antibody for 12 hour at 4°C in PBS + 1% BSA.




Immunocytochemistry - Anti-RAB7 antibody [Rab7-117] - Late Endosome Marker (ab50533)

This image is courtesy of an Abreview submitted by Armen Petrosyan

ab50533 staining RAB7 in HepG2 cells by ICC/IF

(Immunocytochemistry/immunofluorescence). Cells were fixed with formaldehyde, permeabilized with 0.2% Triton X-100 and blocked with 1% serum for 1 hour at 22°C. Samples were incubated with primary antibody (1/50 in PBST + 1% donkey serum) for 3 hours at 22°C. An Alexa Fluor® 488-conjugated donkey anti-mouse IgG polyclonal (1/200) was used as the secondary antibody.



Western blot - Anti-RAB7 antibody [Rab7-117] - Late Endosome Marker (ab50533)

This image is courtesy of an Abreview submitted by Armen Petrosyan

Anti-RAB7 antibody [Rab7-117] - Late Endosome Marker (ab50533) at 1/2000 dilution + Rat hepatocytes whole cell lysate at 20 µg

Secondary

HRP-conjugated donkey anti-mouse IgG polyclonal at 1/10000 dilution

Developed using the ECL technique.


Performed under reducing conditions.

Predicted band size: 23 kDa

Exposure time: 30 seconds

Blocked with 5% milk for 1 hour at 22°C.

Incubated with the primary antibody for 12 hour at 4°C in PBS + 1% BSA.



Western blot - Anti-RAB7 antibody [Rab7-117] - Late Endosome Marker (ab50533)

This image is a courtesy of Anonymous Abreview

Anti-RAB7 antibody [Rab7-117] - Late Endosome Marker (ab50533) at 1/1000 dilution + Lysate prepared from mouse neuroblastoma cells at 10 µg

Secondary

HRP-conjugated goat monoclonal to mouse IgG at 1/2000 dilution


Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 23 kDa

Observed band size: 23 kDa

Exposure time: 20 seconds



Anti-RAB7 antibody [Rab7-117] - Late Endosome Marker (ab50533) at 1/500 dilution (for 3 hours at 20°C) + Human brain whole tissue lysate at 15 µg

Secondary

An HRP-conjugated Goat anti-mouse IgG polyclonal at 1/5000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 23 kDa

Observed band size: 24 kDa

Exposure time: 1 minute

Blocking Step: 5% Milk for 1 hour at 20°C

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors