

Anti-RAP1A antibody [1D2-1C64] ab175329

5 References **5 图像**

概述

产品名称	Anti-RAP1A抗体[1D2-1C64]
描述	小鼠单克隆抗体[1D2-1C64] to RAP1A
宿主	Mouse
经测试应用	适用于: IHC-P, WB, IP, ICC/IF
种属反应性	与反应: Mouse, Human 预测可用于: Rat, Cow 
免疫原	Recombinant full length protein corresponding to Human RAP1A aa 1 to the C-terminus. Database link: P62834 <div>  Run BLAST with  Run BLAST with </div>
阳性对照	HeLa, Hek293T, U2OS and mouse NIH 3T3 cell lysates, HeLa cells, C2C12 cells.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
存储溶液	Preservative: 0.05% Sodium azide Constituents: 30% Glycerol (glycerin, glycerine), 0.1% BSA, 69% PBS
纯度	Protein A purified
克隆	单克隆
克隆编号	1D2-1C64
同种型	IgG2a

应用


The Abpromise guarantee **Abpromise™**承诺保证使用ab175329于以下的经测试应用
“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-P		1/100 - 1/1000.
WB		1/500 - 1/1000. Predicted molecular weight: 21 kDa.
IP		Use at 2 µg/mg of lysate.
ICC/IF		1/50 - 1/200.

靶标

功能	Induces morphological reversion of a cell line transformed by a Ras oncogene. Counteracts the mitogenic function of Ras, at least partly because it can interact with Ras GAPs and RAF in a competitive manner.
序列相似性	Belongs to the small GTPase superfamily. Ras family.
细胞定位	Cell membrane.


图片



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-RAP1A antibody [1D2-1C64] (ab175329)


Immunohistochemistry analysis of RAP1A using ab175329 at 1/200 dilution showing staining in the membrane of paraffin-embedded human kidney tissue (right) compared with a negative control without primary antibody (left). Detection was performed using an HRP-conjugated secondary antibody followed by colorimetric detection using a DAB kit.

Antigen retrieval was performed using 10mM sodium citrate (pH 6.0), microwaved for 8-15 min.



Immunocytochemistry/ Immunofluorescence - Anti-RAP1A antibody [1D2-1C64] (ab175329)

Immunocytochemical analysis of RAP1A (green) showing staining in the cytoplasm of Formalin-fixed and 0.1% Triton X-100 permeabilized C2C12 cells (right) using ab175329 at 1/20 dilution compared to a negative control without primary antibody (left) followed by DyLight-conjugated secondary antibody. F-actin (red) was stained with a fluorescent red phalloidin and nuclei (blue) were stained with DAPI.



Western blot - Anti-RAP1A antibody [1D2-1C64] (ab175329)

All lanes : Anti-RAP1A antibody [1D2-1C64] (ab175329) at 1/500 dilution

Lane 1 : HeLa cell lysate

Lane 2 : HEK293T cell lysate

Lane 3 : U2OS cell lysate

Lane 4 : Mouse NIH 3T3 cell lysate


Lysates/proteins at 25 µg per lane.

Secondary

All lanes : Goat anti-mouse IgG-HRP at 1/15000 dilution


Developed using the ECL technique.

Predicted band size: 21 kDa



Immunocytochemistry/ Immunofluorescence - Anti-RAP1A antibody [1D2-1C64] (ab175329)

Immunofluorescence analysis of formalin-fixed permeabilized HeLa cells, labeling RAP1A (green, left panel) using ab175329 at a 1/100 dilution followed by DyLight 488-conjugated goat anti-mouse IgG secondary antibody at a 1/400 dilution. Nuclei (blue) were stained with Hoechst 33342 dye (central panel).



Immunoprecipitation - Anti-RAP1A antibody [1D2-1C64] (ab175329)

Western blot analysis on immunoprecipitation pellet from mouse NIH 3T3 cells. The antigen-antibody complex was formed by incubating 750 µg of NIH 3T3 whole cell lysate with 2 µg of ab175329 overnight at 4°C. The immune-complex was then captured on 50 µl Protein A/G Plus Agarose, washes extensively and eluted in sample buffer. 1) 25 µg of NIH 3T3 whole cell lysate, as a control, and 2) eluted sample were resolved on a SDS PAGE gel. The membrane was probed with ab175329 at a 1/500 dilution. Chemiluminescent detection was performed.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you

- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors