


Anti-RPL11 antibody ab79352

★★★★★ [3 Abreviews](#) [21 References](#) [4 图像](#)

概述

产品名称	Anti-RPL11抗体
描述	兔多克隆抗体to RPL11
宿主	Rabbit
经测试应用	适用于: IHC-P, WB, IP
种属反应性	与反应: Human, Zebrafish 预测可用于: Mouse, Rat, Cow, Cat, Orangutan 
免疫原	Synthetic peptide corresponding to Human RPL11 aa 150 to the C-terminus (C terminal) conjugated to keyhole limpet haemocyanin. (Peptide available as ab88221)
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	pH: 7.40 Preservative: 0.02% Sodium azide Constituent: PBS Batches of this product that have a concentration < 1mg/ml may have BSA added as a stabilising agent. If you would like information about the formulation of a specific lot, please contact our scientific support team who will be happy to help.
纯度	Immunogen affinity purified
克隆	多克隆

同种型

IgG

应用

The Abpromise guarantee

Abpromise™承诺保证使用ab79352于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-P		Use a concentration of 1 µg/ml. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.
WB	★★★★★ (3)	Use at an assay dependent concentration. Detects a band of approximately 20 kDa (predicted molecular weight: 22 kDa).
IP		Use at an assay dependent concentration.

靶标

功能	Binds to 5S ribosomal RNA (By similarity). Required for rRNA maturation and formation of the 60S ribosomal subunits. Promotes nucleolar location of PML.
疾病相关	Defects in RPL11 are the cause of Diamond-Blackfan anemia type 7 (DBA7) [MIM:612562]. DBA7 is a form of Diamond-Blackfan anemia, a congenital non-regenerative hypoplastic anemia that usually presents early in infancy. Diamond-Blackfan anemia is characterized by a moderate to severe macrocytic anemia, erythroblastopenia, and an increased risk of malignancy. 30 to 40% of Diamond-Blackfan anemia patients present with short stature and congenital anomalies, the most frequent being craniofacial (Pierre-Robin syndrome and cleft palate), thumb and urogenital anomalies.
序列相似性	Belongs to the ribosomal protein L5P family.
细胞定位	Nucleus > nucleolus.

图片



Western blot - Anti-RPL11 antibody (ab79352)

All lanes : Anti-RPL11 antibody (ab79352) at 1 µg/ml

Lane 1 : Human liver tissue lysate - total protein (**ab29889**)

Lane 2 : HepG2 (Human hepatocellular liver carcinoma cell line) Whole Cell Lysate

Lane 3 : Human brain tissue lysate - total protein (**ab29466**)

Lane 4 : HeLa (Human epithelial carcinoma cell line) Whole Cell Lysate

Lane 5 : Jurkat (Human T cell lymphoblast-like cell line) Whole Cell Lysate

Lane 6 : Caco 2 (Human colonic carcinoma cell line) Whole Cell Lysate

Lane 7 : HEK293 (Human embryonic kidney cell line) Whole Cell Lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat polyclonal to Rabbit IgG - H&L - Pre-Adsorbed (HRP) at 1/3000 dilution

Developed using the ECL technique.


Performed under reducing conditions.

Predicted band size: 22 kDa

Observed band size: 22 kDa

Additional bands at: 30 kDa, 45 kDa, 70 kDa. We are unsure as to the identity of these extra bands.

RPL11 contains a number of potential phosphorylation and acetylation sites (SwissProt) which may explain its migration at a higher molecular weight than predicted.




RPL11 was immunoprecipitated using 0.5mg Jurkat whole cell extract, 5µg of Rabbit polyclonal to RPL11 and 50µl of protein G magnetic beads (+). No antibody was added to the control (-). The antibody was incubated under agitation with Protein G beads for 10min, Jurkat whole cell extract lysate diluted in RIPA buffer was added to each sample and incubated for a further 10min under agitation.

Proteins were eluted by addition of 40µl SDS loading buffer and incubated for 10min at 70°C; 10µl of each sample was separated on a SDS PAGE gel, transferred to a nitrocellulose membrane, blocked with 5% BSA and probed with ab79352.

Secondary: Clean blot (HRP conjugate) at 1/1000 dilution.


Band: 22kDa: RPL11



- Anti-RPL11 antibody (ab79352)

IHC image of ab79352 staining in human pancreas formalin fixed paraffin embedded tissue section, performed on a Leica Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab79352, 1µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.



All lanes : Anti-RPL11 antibody (ab79352) at 1 µg/ml

Lane 1 : Marker

Lane 2 : Zebrafish brain homogenate at 20 µg

Lane 3 : Zebrafish heart homogenate at 20 µg

Lane 4 : Zebrafish liver homogenate at 20 µg

Lane 5 : Zebrafish skeletal muscle homogenate at 20 µg

Lane 6 : Human hepatocellular carcinoma cell line at 20 µg

Secondary

All lanes : Goat polyclonal to Rabbit IgG – H&L – Pre-Adsorbed (HRP) at 1/6000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 22 kDa

Observed band size: 22 kDa

Exposure time: 3 minutes

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors