

Anti-RSK1 p90 (phospho S380) antibody [E239] ab32203

重组 RabMAb

13 References [7 图像](#)

概述

产品名称	Anti-RSK1 p90 (phospho S380)抗体[E239]
描述	兔单克隆抗体[E239] to RSK1 p90 (phospho S380)
宿主	Rabbit
特异性	The antibody only detects RSK1 phosphorylated on Serine 380.
经测试应用	适用于: WB, ICC/IF, IP, Dot blot
种属反应性	与反应: Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	IP: C6 whole cell lysate treated with 100nM Calyculin A for 30 minutes. ICC/IF: A431 cells treated with 100ng/ml EGF for 10 minutes. WB: HeLa starved overnight then treated with 200nM TPA for 4 hours whole cell lysate. PC-12 (Rat adrenal gland pheochromocytoma) treated with 200ng/ml NGF for 24 hours whole cell lysate. NIH/3T3 (Mouse embryonic fibroblast) starved for 24 hours then treated with 100nM Calyculin A for 30 minutes whole cell lysate.
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
存储溶液	<p>pH: 7.20</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA</p>
纯度	Protein A purified
克隆	单克隆

克隆编号	E239
同种型	IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab32203于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/10000. Detects a band of approximately 90 kDa (predicted molecular weight: 83 kDa).
ICC/IF		1/50. For the unpurified format use at 1/250 - 1/500 dilution
IP		1/20. For the unpurified format use at 1/80 dilution
Dot blot		1/1000.

靶标

功能	Serine/threonine kinase that may play a role in mediating the growth-factor and stress induced activation of the transcription factor CREB.
序列相似性	<p>Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. S6 kinase subfamily.</p> <p>Contains 1 AGC-kinase C-terminal domain.</p> <p>Contains 2 protein kinase domains.</p>
翻译后修饰	Autophosphorylated on Ser-380, as part of the activation process.

图片



All lanes : Anti-RSK1 p90 (phospho S380) antibody [E239] (ab32203) at 1/1000 dilution

Lane 1 : Untreated PC-12 (Rat adrenal gland pheochromocytoma) whole cell lysate

Lane 2 : PC-12 (Rat adrenal gland pheochromocytoma) treated with 200ng/ml NGF for 24 hours whole cell lysate

Lane 3 : PC-12 (Rat adrenal gland pheochromocytoma) treated with 200ng/ml NGF for 24 hours whole cell lysate, then the membrane treated with Alkaline Phosphatase for 1 hour

Lysates/proteins at 15 µg per lane.

Secondary


All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/20000 dilution

Predicted band size: 83 kDa

Observed band size: 90 kDa

Blocking and dilution buffer: 5% NFDM/TBST.

Image produced using the purified version.



Purified ab32203 at 1:20 dilution (0.8µg) immunoprecipitating RSK1 p90 in C6 whole cell lysate treated with 100nM Calyculin A for 30 minutes.

Lane 1 (input): C6 (Rat glial tumor glial cell) whole cell lysate treated with 100nM Calyculin A for 30 minutes 10µg.

Lane 2 (+): ab32203 + C6 whole cell lysate treated with 100nM Calyculin A for 30 minutes.


Lane 3 (-): Rabbit monoclonal IgG (**ab172730**) instead of ab32203 in C6 whole cell lysate treated with 100nM Calyculin A for 30 minutes.

VeriBlot for IP Detection Reagent (HRP)(**ab131366**) (1:1000 dilution) was used for Western blotting.

Blocking Buffer and concentration: 5% NFDM/TBST.


Diluting buffer and concentration: 5% NFDM/TBST.

Observed band size: 90 kDa



Immunocytochemistry/ Immunofluorescence - Anti-
RSK1 p90 (phospho S380) antibody [E239]
(ab32203)

Immunocytochemistry analysis of A431 (Human epidermoid carcinoma epithelial cell) treated with 100ng/ml EGF for 10 minutes cells labeling RSK1 p90 with Purified ab32203 at 1:50 dilution (2.6 µg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with Ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1:200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, [ab150077](#)) was used as the secondary antibody at 1:1000 (2 µg/ml) dilution. DAPI (blue) was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.




Dot Blot - Anti-RSK1 p90 (phospho S380) antibody
[E239] (ab32203)

This data was produced using the unpurified format.

Dot blot analysis of RSK1 p90 (phospho S380) phospho peptide (Lane 1) and RSK1 p90 non-phospho peptide (Lane 2) labeling RSK1 p90 (phospho S380) with ab32203 at a dilution of 1/1000. [ab97051](#) (Peroxidase conjugated goat anti-rabbit IgG) (H+L) at 1/100 000 was used as the secondary antibody.

Blocking and diluting buffer: 5% NFDm/TBST.

Exposure time: 3 minutes.



Western blot - Anti-RSK1 p90 (phospho S380)
antibody [E239] (ab32203)

All lanes : Anti-RSK1 p90 (phospho S380) antibody [E239]
(ab32203) at 1/1000 dilution

Lane 1 : HeLa (Human cervix adenocarcinoma epithelial cell)
starved overnight whole cell lysate

Lane 2 : HeLa (Human cervix adenocarcinoma epithelial cell)
starved overnight then treated with 200nM TPA for 4 hours whole
cell lysate

Lane 3 : HeLa (Human cervix adenocarcinoma epithelial cell)
starved overnight then treated with 200nM TPA for 4 hours whole
cell lysate, then the membrane treated with Alkaline Phosphatase
for 1 hour

Lysates/proteins at 15 µg per lane.

Secondary


All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

Predicted band size: 83 kDa

Observed band size: 90 kDa

Blocking and dilution buffer: 5% NFDM/TBST.

Image produced using the purified version.



All lanes : Anti-RSK1 p90 (phospho S380) antibody [E239] (ab32203) at 1/1000 dilution

Lane 1 : NIH/3T3 (Mouse embryonic fibroblast) starved for 24 hours whole cell lysate

Lane 2 : NIH/3T3 (Mouse embryonic fibroblast) starved for 24 hours then treated with 100nM Calyculin A for 30 minutes whole cell lysate

Lane 3 : NIH/3T3 (Mouse embryonic fibroblast) starved for 24 hours then treated with 100nM Calyculin A for 30 minutes whole cell lysate, and then the membrane treated with Alkaline Phosphatase for 1 hour

Lysates/proteins at 15 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

Predicted band size: 83 kDa

Observed band size: 90 kDa

Blocking and dilution buffer: 5% NFDM/TBST.

Image produced using the purified version.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-RSK1 p90 (phospho S380) antibody [E239]
(ab32203)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors