

### Anti-S100A8 antibody [EPR3554] ab92331

**重组 RabMAb**

★★★★☆ **7 Abreviews** **43 References** **10 图像**

#### 概述

<b>产品名称</b>	Anti-S100A8抗体[EPR3554]
<b>描述</b>	兔单克隆抗体[EPR3554] to S100A8
<b>宿主</b>	Rabbit
<b>经测试应用</b>	<b>适用于:</b> WB, IP, IHC-P <b>不适用于:</b> Flow Cyt or ICC/IF
<b>种属反应性</b>	<b>与反应:</b> Mouse, Human
<b>免疫原</b>	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
<b>阳性对照</b>	WB: Human (adult and fetal) and mouse spleen tissue lysates and PMBC and HL-60 cell lysates. IHC-P: Human cervical carcinoma, breast carcinoma, tonsil and spleen tissues. IP: HL-60 cell lysate.
<b>常规说明</b>	This product is a recombinant monoclonal antibody, which offers several advantages including: <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> For more information <a href="#">see here</a> . Our RabMAb <sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a> .

#### 性能

<b>形式</b>	Liquid
<b>存放说明</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
<b>存储溶液</b>	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol, 0.05% BSA
<b>纯度</b>	Protein A purified
<b>克隆</b>	单克隆
<b>克隆编号</b>	EPR3554

## 应用

## The Abpromise guarantee

**Abpromise™**承诺保证使用ab92331于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB	★★★★☆ (4)	1/1000. Predicted molecular weight: 11 kDa. <b>For unpurified use at 1/2000 - 1/10000.</b>
IP		1/30. <b>For unpurified use at 1/20.</b>
IHC-P	★★★★★ (1)	1/500 - 1/1000. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. See <a href="#">IHC antigen retrieval protocols</a> . <b>For unpurified use at 1/100 - 1/250.</b>

## 应用说明

Is unsuitable for Flow Cyt or ICC/IF.

## 靶标

## 功能

S100A8 is a calcium- and zinc-binding protein which plays a prominent role in the regulation of inflammatory processes and immune response. It can induce neutrophil chemotaxis and adhesion. Predominantly found as calprotectin (S100A8/A9) which has a wide plethora of intra- and extracellular functions. The intracellular functions include: facilitating leukocyte arachidonic acid trafficking and metabolism, modulation of the tubulin-dependent cytoskeleton during migration of phagocytes and activation of the neutrophilic NADPH-oxidase. Activates NADPH-oxidase by facilitating the enzyme complex assembly at the cell membrane, transferring arachidonic acid, an essential cofactor, to the enzyme complex and S100A8 contributes to the enzyme assembly by directly binding to NCF2/P67PHOX. The extracellular functions involve proinflammatory, antimicrobial, oxidant-scavenging and apoptosis-inducing activities. Its proinflammatory activity includes recruitment of leukocytes, promotion of cytokine and chemokine production, and regulation of leukocyte adhesion and migration. Acts as an alarmin or a danger associated molecular pattern (DAMP) molecule and stimulates innate immune cells via binding to pattern recognition receptors such as Toll-like receptor 4 (TLR4) and receptor for advanced glycation endproducts (AGER). Binding to TLR4 and AGER activates the MAP-kinase and NF-kappa-B signaling pathways resulting in the amplification of the proinflammatory cascade. Has antimicrobial activity towards bacteria and fungi and exerts its antimicrobial activity probably via chelation of Zn(2+) which is essential for microbial growth. Can induce cell death via autophagy and apoptosis and this occurs through the cross-talk of mitochondria and lysosomes via reactive oxygen species (ROS) and the process involves BNIP3. Can regulate neutrophil number and apoptosis by an anti-apoptotic effect; regulates cell survival via ITGAM/ITGB and TLR4 and a signaling mechanism involving MEK-ERK. Its role as an oxidant scavenger has a protective role in preventing exaggerated tissue damage by scavenging oxidants. Can act as a potent amplifier of inflammation in autoimmunity as well as in cancer development and tumor spread. The iNOS-S100A8/A9 transnitrosylase complex directs selective inflammatory stimulus-dependent S-

nitrosylation of GAPDH and probably multiple targets such as ANXA5, EZR, MSN and VIM by recognizing a [IL]-x-C-x-x-[DE] motif; S100A8 seems to contribute to S-nitrosylation site selectivity.

### 组织特异性

Calprotectin (S100A8/9) is predominantly expressed in myeloid cells. Except for inflammatory conditions, the expression is restricted to a specific stage of myeloid differentiation since both proteins are expressed in circulating neutrophils and monocytes but are absent in normal tissue macrophages and lymphocytes. Under chronic inflammatory conditions, such as psoriasis and malignant disorders, also expressed in the epidermis. Found in high concentrations at local sites of inflammation or in the serum of patients with inflammatory diseases such as rheumatoid, cystic fibrosis, inflammatory bowel disease, Crohn's disease, giant cell arteritis, cystic fibrosis, Sjogren's syndrome, systemic lupus erythematosus, and progressive systemic sclerosis. Involved in the formation and deposition of amyloids in the aging prostate known as corpora amylacea inclusions. Strongly up-regulated in many tumors, including gastric, esophageal, colon, pancreatic, bladder, ovarian, thyroid, breast and skin cancers.


### 序列相似性

Belongs to the S-100 family.  
Contains 2 EF-hand domains.

### 细胞定位

Secreted. Cytoplasm. Cytoplasm, cytoskeleton. Cell membrane. Predominantly localized in the cytoplasm. Upon elevation of the intracellular calcium level, translocated from the cytoplasm to the cytoskeleton and the cell membrane. Upon neutrophil activation or endothelial adhesion of monocytes, is secreted via a microtubule-mediated, alternative pathway.

### 图片



**All lanes :** Anti-S100A8 antibody [EPR3554] (ab92331) at 1/20000 dilution (purified)

**Lane 1 :** HL-60 (Human promyelocytic leukemia cell line) cell lysate

**Lane 2 :** Human fetal spleen tissue lysate

Lysates/proteins at 20 µg per lane.


#### Secondary

**All lanes :** Peroxidase-conjugated goat anti-rabbit IgG (H+L) at 1/1000 dilution

**Predicted band size:** 11 kDa

**Observed band size:** 11 kDa

**Blocking/Dilution buffer:** 5% NFDm/TBST.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-S100A8 antibody [EPR3554] (ab92331)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human cervix carcinoma tissue labeling S100A8 with purified ab92331 at 1/500.

Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9.0. **ab97051**, an HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500).

**Negative control** using PBS instead of primary antibody. Counterstained with hematoxylin.




Immunoprecipitation - Anti-S100A8 antibody [EPR3554] (ab92331)

ab92331 (purified) at 1/30 immunoprecipitating S100A8 in HL-60 (Human promyelocytic leukemia cell line) cell lysate (**Lane 1**).

**Lane 2** - rabbit monoclonal IgG instead of ab92331 in HL-60 lysates. For western blotting, a peroxidase-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/1500).

**Blocking/Dilution buffer:** 5% NFDm/TBST.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-S100A8 antibody [EPR3554] (ab92331)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human cervical carcinoma tissue labeling S100A8 with unpurified ab92331.

**Detection:** DAB staining.

Heat mediated antigen retrieval was performed via the pressure cooker method before commencing with IHC staining protocol.



Western blot - Anti-S100A8 antibody [EPR3554] (ab92331)

Anti-S100A8 antibody [EPR3554] (ab92331) at 1/2000 dilution (purified) + Mouse spleen tissue lysate at 20  $\mu$ g


**Secondary**

Peroxidase-conjugated goat anti-rabbit IgG (H+L) at 1/1000 dilution

**Predicted band size:** 11 kDa


**Observed band size:** 11 kDa

**Blocking/Dilution buffer:** 5% NFDM/TBST.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-S100A8 antibody [EPR3554] (ab92331)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human breast carcinoma tissue labeling S100A8 with unpurified ab92331. Heat mediated antigen retrieval was performed via the pressure cooker method before commencing with IHC staining protocol.



Western blot - Anti-S100A8 antibody [EPR3554] (ab92331)

**All lanes :** Anti-S100A8 antibody [EPR3554] (ab92331) (unpurified)

**Lane 1 :** Human spleen tissue lysate

**Lane 2 :** PBMC cell lysate


**Lane 3 :** HL-60 (Human promyelocytic leukemia cell line) cell lysate

Lysates/proteins at 10 µg per lane.

**Secondary**

**All lanes :** HRP-conjugated goat anti-rabbit IgG at 1/2000 dilution

**Predicted band size:** 11 kDa




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-S100A8 antibody [EPR3554] (ab92331)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human spleen tissue labeling S100A8 with unpurified ab92331.

**Detection:** DAB staining.

Heat mediated antigen retrieval was performed via the pressure cooker method before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-S100A8 antibody [EPR3554] (ab92331)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human tonsil tissue labeling S100A8 with unpurified ab92331. Heat mediated antigen retrieval was performed via the pressure cooker method before commencing with IHC staining protocol.

### Why choose a recombinant antibody?



**Research with confidence**  
Consistent and reproducible results



**Long-term and scalable supply**  
Recombinant technology



**Success from the first experiment**  
Confirmed specificity



**Ethical standards compliant**  
Animal-free production

Anti-S100A8 antibody [EPR3554] (ab92331)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors