

Anti-Smac/Diablo antibody [SMAC 17 1-87] ab111893

1 References **8 图像**

概述

产品名称	Anti-Smac/Diablo抗体[SMAC 17 1-87]
描述	小鼠单克隆抗体[SMAC 17 1-87] to Smac/Diablo
宿主	Mouse
经测试应用	适用于: ICC/IF, IHC-P, WB
种属反应性	与反应: Mouse, Rat, Human
免疫原	Recombinant full length Human Smac/ Diablo with a His tag at C terminal end.
阳性对照	IHC: human colon and testis tissue, mouse testis tissue; ICC/IF: A431, PC-12, MCF-7; WB: PC-12, HeLa, 293T cells, NIH3T3, MCF-7, Hep G2, Jurkat
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid repeated freeze / thaw cycles.
存储溶液	Preservative: 0.05% Sodium azide Constituents: 99% PBS, 0.1% BSA
纯度	Protein G purified
克隆	单克隆
克隆编号	SMAC 17 1-87
同种型	IgG1

应用

The Abpromise guarantee

Abpromise™ 承诺保证使用ab111893于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC/IF		1/20 - 1/200.
IHC-P		1/20 - 1/200. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
WB		Use at an assay dependent concentration. Detects a band of approximately 25 kDa (predicted molecular weight: 27 kDa).

靶标

功能	Promotes apoptosis by activating caspases in the cytochrome c/Apaf-1/caspase-9 pathway. Acts by opposing the inhibitory activity of inhibitor of apoptosis proteins (IAP).
组织特异性	Ubiquitously expressed with highest expression in testis. Expression is also high in heart, liver, kidney, spleen, prostate and ovary. Low in brain, lung, thymus and peripheral blood leukocytes.
结构域	The mature N-terminus mediates interaction with XIAP.
细胞定位	Mitochondrion. Released into the cytosol when cells undergo apoptosis.

图片



All lanes : Anti-Smac/Diablo antibody [SMAC 17 1-87] (ab111893) at 1 µg/ml

Lane 1 : HeLa cell lysate

Lane 2 : Jurkat cell lysate

Lane 3 : MCF-7 cell lysate

Lane 4 : Hep G2 cell lysate

Lane 5 : NIH/3T3 cell lysate


Lane 6 : PC-12 cell lysate

Lysates/proteins at 30 µg per lane.

Secondary


All lanes : Goat anti-Mouse IgG (H+L) Superclonal™ Secondary Antibody, HRP conjugate at 1/4000 dilution

Predicted band size: 27 kDa




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Smac/Diablo antibody [SMAC 17 1-87] (ab111893)

Immunohistochemistry (paraffin-treated sections) analysis of human colon tissue labelling SMAC/Diablo with ab111893 at 1/100 dilution. Tissues were counterstained with hematoxylin and dehydrated with xylene and ethanol. Negative control (left) has no primary antibody. Heat mediated antigen retrieval performed with citrate buffer pH 6 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Smac/Diablo antibody [SMAC 17 1-87] (ab111893)

Immunohistochemistry (paraffin-treated sections) analysis of mouse testis tissue labelling SMAC/Diablo with ab111893 at 1/100 dilution. Tissues were counterstained with hematoxylin and dehydrated with xylene and ethanol. Negative control (left) has no primary antibody. Heat mediated antigen retrieval performed with citrate buffer pH 6 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Smac/Diablo antibody [SMAC 17 1-87] (ab111893)

Immunohistochemistry (paraffin-treated sections) analysis of human testis tissue labelling SMAC/Diablo with ab111893 at 1/100 dilution. Tissues were counterstained with hematoxylin and dehydrated with xylene and ethanol. Negative control (left) has no primary antibody. Heat mediated antigen retrieval performed with citrate buffer pH 6 before commencing with IHC staining protocol.




Immunocytochemistry/ Immunofluorescence - Anti-Smac/Diablo antibody [SMAC 17 1-87] (ab111893)

ab111893 labelling Smac / Diablo (green) in the cytoplasm of A431 cells by Immunocytochemistry/Immunofluorescence. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were incubated with the primary antibody (1:100 in 3% BSA-PBS) overnight at 4 °C. A DyLight-conjugated anti-mouse IgG was used as the secondary antibody. Red (phalloidin) - F-actin, Blue - nuclei. Images were taken at a magnification of 60x.




Immunocytochemistry/ Immunofluorescence - Anti-Smac/Diablo antibody [SMAC 17 1-87] (ab111893)

ab111893 labelling Smac / Diablo (green) in the cytoplasm of MCF-7 cells by Immunocytochemistry/Immunofluorescence. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were incubated with the primary antibody (1:100 in 3% BSA-PBS) overnight at 4 °C. A DyLight-conjugated anti-mouse IgG was used as the secondary antibody. Red (phalloidin) - F-actin, Blue - nuclei. Images were taken at a magnification of 60x.



Immunocytochemistry/ Immunofluorescence - Anti-Smac/Diablo antibody [SMAC 17 1-87] (ab111893)

ab111893 labelling Smac / Diablo (green) in the cytoplasm of PC12 cells by Immunocytochemistry/Immunofluorescence. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were incubated with the primary antibody (1:100 in 3% BSA-PBS) overnight at 4 °C. A DyLight-conjugated anti-mouse IgG was used as the secondary antibody. Red (phalloidin) - F-actin, Blue - nuclei. Images were taken at a magnification of 60x.



All lanes : Anti-Smac/Diablo antibody [SMAC 17 1-87] (ab111893) at 1/1000 dilution

Lane 1 : 293 cell lysate

Lane 2 : HeLa cell lysate

Lane 3 : PC12 cell lysate

Lysates/proteins at 25 µg per lane.

Predicted band size: 27 kDa

Observed band size: 21 kDa

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors