

### Anti-Smad2 antibody [EP567Y] - BSA and Azide free ab216454

敲除验证
重组
RabMAb

[6 References](#)
[6 图像](#)

#### 概述

产品名称	Anti-Smad2抗体[EP567Y] - BSA and Azide free
描述	兔单克隆抗体[EP567Y] to Smad2 - BSA and Azide free
宿主	Rabbit
特异性	This antibody detects a region about 40AA before the MH2 region (not the MH2 region itself).
经测试应用	<b>适用于:</b> ChIC/CUT&RUN-seq, Flow Cyt (Intra), WB, ICC/IF <b>不适用于:</b> IHC-P or IP
种属反应性	<b>与反应:</b> Mouse, Human <b>预测可用于:</b> Rat 
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: A549, HeLa and Jurkat cell lysates. ICC/IF: A673 cells. Flow Cyt (intra): Jurkat and PC3 cells. ChIC/CUT&RUN seq: HaCaT cell.
常规说明	<p>ab216454 is the carrier-free version of <a href="#">ab33875</a>.</p> <p>Our <b>carrier-free</b> antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our <b>conjugation kits</b> for antibody conjugates that are ready-to-use in as little as 20 minutes with &lt;1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar<sup>®</sup> Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar<sup>®</sup> is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> <p>For more information <a href="#">see here</a>.</p>

Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to [RabMAb<sup>®</sup> patents](#).

## 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C. Do Not Freeze.
存储溶液	pH: 7.20 Constituent: PBS
无载体	是
纯度	Protein A purified
克隆	单克隆
克隆编号	EP567Y
同种型	IgG

## 应用

**The Abpromise guarantee**      **Abpromise<sup>™</sup>承诺保证使用ab216454于以下的经测试应用**

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ChIC/CUT&RUN-seq		Use at an assay dependent concentration.
Flow Cyt (Intra)		Use at an assay dependent concentration. <b>ab199376</b> - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
WB		Use at an assay dependent concentration. Detects a band of approximately 58 kDa (predicted molecular weight: 58 kDa).
ICC/IF		Use at an assay dependent concentration.

**应用说明**      Is unsuitable for IHC-P or IP.

## 靶标

功能	Receptor-regulated SMAD (R-SMAD) that is an intracellular signal transducer and transcriptional modulator activated by TGF-beta (transforming growth factor) and activin type 1 receptor kinases. Binds the TRE element in the promoter region of many genes that are regulated by TGF-beta and, on formation of the SMAD2/SMAD4 complex, activates transcription. May act as a tumor suppressor in colorectal carcinoma.
组织特异性	Expressed at high levels in skeletal muscle, heart and placenta.
序列相似性	Belongs to the dwarfin/SMAD family. Contains 1 MH1 (MAD homology 1) domain. Contains 1 MH2 (MAD homology 2) domain.


翻译后修饰

Phosphorylated on one or several of Thr-220, Ser-245, Ser-250, and Ser-255. In response to TGF-beta, phosphorylated on Ser-465/467 by TGF-beta and activin type 1 receptor kinases. Able to interact with SMURF2 when phosphorylated on Ser-465/467, recruiting other proteins, such as SNON, for degradation. In response to decorin, the naturally occurring inhibitor of TGF-beta signaling, phosphorylated on Ser-240 by CaMK2. Phosphorylated by MAPK3 upon EGF stimulation; which increases transcriptional activity and stability, and is blocked by calmodulin. In response to TGF-beta, ubiquitinated by NEDD4L; which promotes its degradation. Acetylated on Lys-19 by coactivators in response to TGF-beta signaling, which increases transcriptional activity. Isoform short: Acetylation increases DNA binding activity in vitro and enhances its association with target promoters in vivo. Acetylation in the nucleus by EP300 is enhanced by TGF-beta.

细胞定位

Cytoplasm. Nucleus. Cytoplasmic and nuclear in the absence of TGF-beta. On TGF-beta stimulation, migrates to the nucleus when complexed with SMAD4. On dephosphorylation by phosphatase PPM1A, released from the SMAD2/SMAD4 complex, and exported out of the nucleus by interaction with RANBP1.


图片



This data was developed using the same antibody clone in a different buffer formulation ([ab33875](#)).

ChIC/CUT&RUN was performed using a pAG-MNase at a final concentration of 700 ng/μL, 2.5 x 10<sup>5</sup> HaCaT (Human keratinocyte cell line) cells (treated with 7ng/ml TGF-β for 1h) and 5 μg of [ab33875](#) [EP567Y]. The resulting DNA was sequenced on the Illumina NovaSeq 6000 to a depth of 10 million reads. The negative IgG control [ab172730](#) is also shown.

Additional screenshots of mapped reads can be downloaded [here](#). The University of Geneva owns patents relevant to ChIC (Chromatin Immuno-Cleavage) methods.



Western blot - Anti-Smad2 antibody [EP567Y] - BSA and Azide free (ab216454)

**All lanes :** Anti-Smad2 antibody [EP567Y] ([ab33875](#)) at 1/1000 dilution

**Lane 1 :** Wild-type A549 cell lysate

**Lane 2 :** Jurkat cell lysate

**Lane 3 :** Wild-type HeLa cell lysate

**Lane 4 :** SMAD2 knockout HeLa cell lysate

Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.


**Predicted band size:** 58 kDa

**Observed band size:** 58 kDa

This data was developed using the same antibody clone in a different buffer formulation ([ab33875](#)).

**Lanes 1 -4:** Merged signal (red and green). Green - [ab33875](#) observed at 58 kDa. Red - loading control, [ab8245](#) observed at 37 kDa.


[ab33875](#) was shown to react with Smad2 in wild-type HeLa. Loss of signal was observed when knockout cell line [ab255430](#) (knockout cell lysate [ab263833](#)) was used. Wild-type and Smad2 knockout samples were subjected to SDS-PAGE. [ab33875](#) and Anti-GAPDH antibody [6C5] - Loading Control ([ab8245](#)) were incubated overnight at 4°C at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed ([ab216776](#)) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Flow Cytometry (Intracellular) - Anti-Smad2 antibody  
[EP567Y] - BSA and Azide free (ab216454)

Overlay histogram showing Jurkat cells stained with purified **ab33875** (pink line) at a dilution of 1/110. The cells were fixed with 2% PFA. FITC goat anti-rabbit was used at a dilution of 1/150 and rabbit monoclonal IgG was used as the isotype control (green line).


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab33875**).



Immunocytochemistry/ Immunofluorescence - Anti-Smad2 antibody [EP567Y] - BSA and Azide free (ab216454)

Immunofluorescent staining of A673 cells, fixed with 4% PFA, using purified **ab33875** at a dilution of 1/300. An Alexa Fluor<sup>®</sup> 555 goat anti-rabbit was used at 1/200. The negative control is shown in the bottom right hand panel - for the negative control, purified **ab33875** was used at a dilution of 1/200 followed by an Alexa Fluor<sup>®</sup> 555 goat anti-mouse antibody at a dilution of 1/500.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab33875**).



Flow Cytometry (Intracellular) - Anti-Smad2 antibody  
[EP567Y] - BSA and Azide free (ab216454)

Overlay histogram showing PC3 cells stained with unpurified **ab33875** (red line). The cells were fixed with methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (**ab33875**, 1/100 dilution) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-rabbit IgG (H+L) (**ab96899**) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit monoclonal IgG (1 µg/1x10<sup>6</sup> cells) used under the same conditions. Acquisition of >5,000 events was performed. This antibody gave a decreased signal in PC3 cells fixed with 4% paraformaldehyde (10 min)/permeabilized in 0.1% PBS-Tween used under the same conditions.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab33875**).

#### Why choose a recombinant antibody?



**Research with confidence**  
Consistent and reproducible results



**Long-term and scalable supply**  
Recombinant technology



**Success from the first experiment**  
Confirmed specificity



**Ethical standards compliant**  
Animal-free production

Anti-Smad2 antibody [EP567Y] - BSA and Azide free (ab216454)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish

- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

#### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors