

### Anti-Smad2 antibody [EP784Y] ab40855

敲除验证
重组
RabMAb

★★★★★
[4 Abreviews](#)
[89 References](#)
[14 图像](#)

#### 概述

产品名称	Anti-Smad2抗体[EP784Y]
描述	兔单克隆抗体[EP784Y] to Smad2
宿主	Rabbit
特异性	This antibody is specific for MH 1 domain of Smad2.
经测试应用	<b>适用于:</b> IHC-P, IP, ICC/IF, Flow Cyt (Intra), ChIC/CUT&RUN-seq, WB
种属反应性	<b>与反应:</b> Rat, Human
免疫原	Synthetic peptide within Human Smad2 aa 50-150. The exact sequence is proprietary.
阳性对照	WB: A549, Jurkat, HeLa, A-673, HUVEC and C6 cell lysates. IP: HeLa IHC-P: Human bladder and prostate carcinoma tissue. ICC/IF: HeLa cells. Flow Cyt (intra): HeLa and PC3 cells. ChIC/CUT&RUN seq: HaCaT cell
常规说明	<p><b>The rat recommendation is based on the WB results. This antibody may not be suitable for IHC with rat samples.</b></p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> <p>For more information <a href="#">see here</a>.</p> <p>Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a>.</p> <p>Mouse: We have preliminary internal testing data to indicate this antibody may not react with this species. Please contact us for more information.</p>

#### 性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
存储溶液	<p>pH: 7.20</p> <p>Preservative: 0.01% Sodium azide</p>

	Constituents: 40% Glycerol, 0.05% BSA, 59% PBS
纯度	Protein A purified
克隆	单克隆
克隆编号	EP784Y
同种型	IgG

应用

The Abpromise guarantee      **Abpromise™**承诺保证使用ab40855于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-P		1/50. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
IP		1/20 - 1/50.
ICC/IF		1/100 - 1/250.
Flow Cyt (Intra)		1/20 - 1/100. <b>ab172730</b> - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
ChIC/CUT&RUN-seq		Use at an assay dependent concentration.
WB	★★★★★ (4)	1/2000 - 1/10000. Detects a band of approximately 55 kDa (predicted molecular weight: 58 kDa).

靶标


功能	Receptor-regulated SMAD (R-SMAD) that is an intracellular signal transducer and transcriptional modulator activated by TGF-beta (transforming growth factor) and activin type 1 receptor kinases. Binds the TRE element in the promoter region of many genes that are regulated by TGF-beta and, on formation of the SMAD2/SMAD4 complex, activates transcription. May act as a tumor suppressor in colorectal carcinoma.
组织特异性	Expressed at high levels in skeletal muscle, heart and placenta.
序列相似性	Belongs to the dwarfin/SMAD family. Contains 1 MH1 (MAD homology 1) domain. Contains 1 MH2 (MAD homology 2) domain.
翻译后修饰	Phosphorylated on one or several of Thr-220, Ser-245, Ser-250, and Ser-255. In response to TGF-beta, phosphorylated on Ser-465/467 by TGF-beta and activin type 1 receptor kinases. Able to interact with SMURF2 when phosphorylated on Ser-465/467, recruiting other proteins, such as SNON, for degradation. In response to decorin, the naturally occurring inhibitor of TGF-beta signaling, phosphorylated on Ser-240 by CaMK2. Phosphorylated by MAPK3 upon EGF stimulation; which increases transcriptional activity and stability, and is blocked by calmodulin. In response to TGF-beta, ubiquitinated by NEDD4L; which promotes its degradation. Acetylated on Lys-19 by coactivators in response to TGF-beta signaling, which increases

transcriptional activity. Isoform short: Acetylation increases DNA binding activity in vitro and enhances its association with target promoters in vivo. Acetylation in the nucleus by EP300 is enhanced by TGF-beta.

## 细胞定位

Cytoplasm. Nucleus. Cytoplasmic and nuclear in the absence of TGF-beta. On TGF-beta stimulation, migrates to the nucleus when complexed with SMAD4. On dephosphorylation by phosphatase PPM1A, released from the SMAD2/SMAD4 complex, and exported out of the nucleus by interaction with RANBP1.


## 图片



ChIC/CUT&RUN sequencing - Anti-Smad2 antibody [EP784Y] (ab40855)

ChIC/CUT&RUN was performed using a pAG-MNase at a final concentration of 700 ng/ $\mu$ L,  $2.5 \times 10^5$  HaCaT (Human keratinocyte cell line) cells (treated with 7ng/ml TGF- $\beta$  for 1h) and 5  $\mu$ g of ab40855 [EP784Y]. The resulting DNA was sequenced on the Illumina NovaSeq 6000 to a depth of 10 million reads. The negative IgG control **ab172730** is also shown.

Additional screenshots of mapped reads can be downloaded [here](#). The University of Geneva owns patents relevant to ChIC (Chromatin Immuno-Cleavage) methods.



Western blot - Anti-Smad2 antibody [EP784Y] (ab40855)

**All lanes** : Anti-Smad2 antibody [EP784Y] (ab40855) at 1/2000 dilution

**Lane 1** : A549 cell lysate

**Lane 2** : Jurkat cell lysate

**Lane 3** : Wild-type HeLa cell lysate

**Lane 4** : SMAD2 knockout HeLa cell lysate


Lysates/proteins at 20  $\mu$ g per lane.

**Predicted band size:** 58 kDa

**Lanes 1 - 4:** Merged signal (red and green). Green - ab40855 observed at 58 kDa. Red - loading control, **ab8245** observed at 37 kDa.

ab40855 was shown to react with Smad2 in wild-type HeLa. Loss of signal was observed when knockout cell line **ab255430**

(knockout cell lysate [ab263833](#)) was used. Wild-type and Smad2 knockout samples were subjected to SDS-PAGE. ab40855 and Anti-GAPDH antibody [6C5] - Loading Control ([ab8245](#)) were incubated overnight at 4°C at 1 in 2000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed ([ab216776](#)) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



**All lanes :** Anti-Smad2 antibody [EP784Y] (ab40855) at 1/1000 dilution

**Lane 1 :** Wild-type HeLa whole cell lysate

**Lane 2 :** Smad2 knockout HeLa whole cell lysate


**Lane 3 :** A549 whole cell lysate

Lysates/proteins at 20 µg per lane.

**Predicted band size:** 58 kDa

**Lanes 1 - 3:** Merged signal (red and green). Green - ab40855 observed at 58 kDa. Red - loading control, [ab9484](#), observed at 37 kDa.

ab40855 was shown to specifically react with Smad2 in wild-type HeLa cells as signal was lost in Smad2 knockout cells. Wild-type and SMAD2 knockout samples were subjected to SDS-PAGE. Ab40855 and [ab9484](#) (Mouse anti-GAPDH loading control) were incubated overnight at 4°C at 1/10000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed [ab216773](#) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed [ab216776](#) secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.



**All lanes :** Anti-Smad2 antibody [EP784Y] (ab40855) at 1/2000 dilution

**Lane 1 :** A-673 (Human muscle Ewing's Sarcoma cell line) whole cell lysate

**Lane 2 :** HUVEC (Human umbilical vein endothelial cell line) whole cell lysate

**Lane 3 :** C6 (Rat glial tumor cell line) whole cell lysate


Lysates/proteins at 20 µg per lane.

### Secondary

**All lanes :** Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/20000 dilution (Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated)


**Predicted band size:** 58 kDa

Diluting and blocking buffer: 5% NFDM /TBST



ab40855 staining Smad2 in HeLa (human cervix adenocarcinoma) cells by ICC/IF

(Immunocytochemistry/immunofluorescence). Cells were fixed with 4% Paraformaldehyde and permeabilized with 0.1% Triton X-100. Samples were incubated with primary antibody at a dilution of 1/100. A goat anti rabbit IgG (Alexa Fluor® 488) (**ab150077**) was used as the secondary antibody at a dilution of 1/1000. **ab195889** was used as a counterstain for primary antibody ab40855 at 1/1000. DAPI was used as a nuclear counterstain and PBS as a negative control.



**All lanes :** Anti-Smad2 antibody [EP784Y] (ab40855) at 1/10000 dilution

**Lane 1 :** Jurkat (Human T cell leukemia cell line from peripheral blood) whole cell lysate

**Lanes 2-3 :** HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate


Lysates/proteins at 20 µg per lane.

### Secondary

**All lanes :** Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 20000 µg (Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated)


**Predicted band size:** 58 kDa

Blocking and diluting buffer: 5% NFDM/TBST



ab40855 staining Smad2 in human bladder carcinoma tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with paraformaldehyde and antigen retrieval was by heat mediation in a EDTA buffer. Samples were incubated with primary antibody at a dilution of 1/50. A ImmunoHistoProbe one step HRP Polymer was used as a secondary antibody, ready to use.

**Negative control 1:** PBS in place of primary antibody.



Immunoprecipitation - Anti-Smad2 antibody  
[EP784Y] (ab40855)

ab40855 (purified) at 1/20 immunoprecipitating EGFR in HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate.


Lane 1 (input): HeLa whole cell lysate (10µg)

Lane 2 (+): ab40855 + HeLa whole cell lysate.

Lane 3 (-): Rabbit monoclonal IgG (**ab172730**) instead of ab40855 in HeLa whole cell lysate.

For western blotting, **ab131366** VeriBlot for IP (HRP) was used for detection (1/1000).

Blocking/Diluting buffer and concentration: 5% NFDm/TBST.




Flow Cytometry (Intracellular) - Anti-Smad2 antibody  
[EP784Y] (ab40855)

ab40855 staining Smad2 in the human cell line HeLa (Human epithelial cell line from cervix adenocarcinoma) by intracellular flow cytometry. Cells were fixed with 4% paraformaldehyde and the sample was incubated with the primary antibody at a dilution of 1/20. A goat anti rabbit IgG (Alexa Fluor® 488) at a dilution of 1/2000 was used as the secondary antibody.


Isoytype control: Rabbit monoclonal IgG (Black)

Unlabelled control: Cell without incubation with primary antibody and secondary antibody (Blue)



Immunocytochemistry/ Immunofluorescence - Anti-Smad2 antibody [EP784Y] (ab40855)

Immunofluorescence staining of HeLa cells with purified ab40855 at a working dilution of 1/500, counter-stained with DAPI. The secondary antibody was an Alexa Fluor<sup>®</sup> 488 conjugated goat anti-rabbit (**ab150077**), used at a dilution of 1/1000. The cells were fixed in 4% PFA and permeabilized using 0.1% Triton X 100. The negative control is shown in bottom right hand panel - for the negative control, PBS was used instead of the primary antibody.



Flow Cytometry (Intracellular) - Anti-Smad2 antibody [EP784Y] (ab40855)

Overlay histogram showing PC3 cells stained with ab40855 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab40855, 1/100 dilution) for 30 min at 22°C. The secondary antibody used was Alexa Fluor<sup>®</sup> 488 goat anti-rabbit IgG (H&L) (**ab150077**) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (1µg/1x10<sup>6</sup> cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter.






Anti-Smad2 antibody [EP784Y] (ab40855) at 1/500000 dilution + Jurkat cell lysate

**Predicted band size:** 58 kDa

**Observed band size:** 58 kDa

Western blot - Anti-Smad2 antibody [EP784Y] (ab40855)



ab40855 at a 1:100 dilution staining Smad2 in human prostate carcinoma tissue.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Smad2 antibody [EP784Y] (ab40855)

### Why choose a recombinant antibody?



**Research with confidence**  
Consistent and reproducible results



**Long-term and scalable supply**  
Recombinant technology



**Success from the first experiment**  
Confirmed specificity



**Ethical standards compliant**  
Animal-free production

Anti-Smad2 antibody [EP784Y] (ab40855)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors