abcam

Product datasheet

Anti-SOCS2 antibody [EPR2588(2)] ab109245

重组 RabMAb

<u>4 References</u> 5 图像

概述			
产 品名称	Anti-SOCS2抗体[EPR2588(2)]		
描述	兔单克隆抗体[EPR2588(2)] to SOCS2		
宿主	Rabbit		
经测试应 用	适用于: WB, IP, Flow Cyt (Intra), ICC/IF		
种属反 应性	与反应: Human		
	预测可用于: Mouse, Rat 🛛 🔺		
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.		
阳性 对 照	WB: NCCIT and K-562 whole cell lysates; IP: K-562 whole cell lysate; Flow Cyt (intra): HeLa cells; ICC/IF: MCF7 cells.		
常规说明	 This product is a recombinant monoclonal antibody, which offers several advantages including: High batch-to-batch consistency and reproducibility Improved sensitivity and specificity Long-term security of supply Animal-free production For more information <u>see here</u>. Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <u>RabMAb[®] patents</u>. 		
性能			
形式	Liquid		
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Stable for 12 months at -20°C.		
存储溶液	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 0.05% BSA, 40% Glycerol (glycerin, glycerine), 59% PBS		
纯 度	Protein A purified		

克隆 单克隆

克隆编号 EPR2588(2)

应用

The Abpromise guarantee Abpromise™承诺保证使用ab109245于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/1000 - 1/10000. Predicted molecular weight: 22 kDa.
IP		1/10 - 1/100.
Flow Cyt (Intra)		1/150.
ICC/IF		1/100 - 1/250.

靶标

功能	SOCS family proteins form part of a classical negative feedback system that regulates cytokine signal transduction. SOCS2 appears to be a negative regulator in the growth hormone/IGF1 signaling pathway. Probable substrate recognition component of a SCF-like ECS (Elongin BC-CUL2/5-SOCS-box protein) E3 ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins.
组织 特异性	High expression in heart, placenta, lung, kidney and prostate.
通路	Protein modification; protein ubiquitination.
序列相似性	Contains 1 SH2 domain. Contains 1 SOCS box domain.
结 构域	The SOCS box domain mediates the interaction with the Elongin BC complex, an adapter module in different E3 ubiquitin ligase complexes.

图片



Western blot - Anti-SOCS2 antibody [EPR2588(2)] (ab109245) **All lanes :** Anti-SOCS2 antibody [EPR2588(2)] (ab109245) at 1/2000 dilution (purified)


Lane 1 : NCCIT (Human pluripotent embryonic carcinoma epithelial cell) whole cell lysate
Lane 2 : K-562 (Human chronic myelogenous leukemia lymphoblast) whole cell lysate

Lysates/proteins at 20 µg per lane.


Secondary

All lanes : Goat Anti-Rabbit IgG (HRP) with minimal cross-reactivity with human IgG at 1/2000 dilution

Predicted band size: 22 kDa Observed band size: 22 kDa




Immunocytochemistry/ Immunofluorescence - Anti-SOCS2 antibody [EPR2588(2)] (ab109245) Immunocytochemistry/ Immunofluorescence analysis of MCF7 (Human breast adenocarcinoma epithelial cell) cells labeling SOCS2 with purified ab109245 at 1/100 dilution (10 μ g/mL). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% TritonX-100. Cells were counterstained with Ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor[®] 594) 1/200 (2.5 μ g/mL). Goat anti rabbit lgG (Alexa Fluor[®] 488, **ab150077**) was used as the secondary antibody at 1/1000 (2 μ g/mL) dilution. DAPI (blue) was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Intracellular Flow Cytometry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling SOCS2 with purified ab109245 at 1/150 dilution (10µg/mL) (Red). Cells were fixed with 4% Paraformaldehyde and permeabilised with 90% Methanol. A Goat anti rabbit IgG (Alexa Fluorr[®] 488, **ab150077**) secondary antibody was used at 1/2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).

Flow Cytometry (Intracellular) - Anti-SOCS2 antibody [EPR2588(2)] (ab109245)



Immunoprecipitation - Anti-SOCS2 antibody [EPR2588(2)] (ab109245)

> Why choose a recombinant antibody? Research with Long-term and confidence scalable supply Consistent and Recombinant reproducible results technology Success from the Ethical standards first experiment compliant Confirmed Animal-free specificity production

Anti-SOCS2 antibody [EPR2588(2)] (ab109245)

Purified ab109245 at 1/70 dilution (2 μg) immunoprecipitating SOCS2 in K-562 whole cell lysate. Lane 1 (input): K-562 (Human chronic myelogenous leukemia lymphoblast) whole cell lysate 10 μg Lane 2 (+): ab109245 + K-562 whole cell lysate. Lane 3 (-): Rabbit monoclonal lgG (**ab172730**) instead of ab109245 in K-562 whole cell lysate. VeriBlot for IP Detection Reagent (HRP) (**ab131366**) (1/1000 dilution) was used for Western blotting. Blocking Buffer and concentration: 5% NFDM/TBST. Diluting buffer and concentration: 5% NFDM/TBST.

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <u>https://www.abcam.cn/abpromise</u> or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors