

# Anti-Sumo 1 antibody [Y299] - BSA and Azide free ab219724

敲除验证
重组
RabMAb

[1 References](#)
[15 图像](#)

### 概述

<b>产品名称</b>	Anti-Sumo 1抗体[Y299] - BSA and Azide free
<b>描述</b>	兔单克隆抗体[Y299] to Sumo 1 - BSA and Azide free
<b>宿主</b>	Rabbit
<b>特异性</b>	This antibody recognises small ubiquitin-related modifier-1 (SUMO-1), also known as SMT3, Sentrin, GMP1 UBL1 and PIC1.
<b>经测试应用</b>	<b>适用于:</b> ICC/IF, Flow Cyt (Intra), IP, WB, IHC-P, ChIP
<b>种属反应性</b>	<b>与反应:</b> Mouse, Rat, Human
<b>免疫原</b>	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
<b>阳性对照</b>	WB: HeLa, A549, C6 and NIH/3T3 cell lysates. Wild-type HAP1 whole cell lysate. IHC-P: Human endometrium, lung carcinoma and bladder carcinoma tissue. Rat stomach tissue. Mouse kidney tissue. ICC/IF: HeLa cells. Flow Cyt (intra): HeLa cells. ChIP: Chromatin prepared from SK-OV-3 cells. IP: NIH/3T3 cell lysate.
<b>常规说明</b>	<p>ab219724 is the carrier-free version of <a href="#">ab32058</a>.</p> <p>Our <b>carrier-free</b> antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our <b>conjugation kits</b> for antibody conjugates that are ready-to-use in as little as 20 minutes with &lt;1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar<sup>®</sup> Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar<sup>®</sup> is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> <p>For more information <a href="#">see here</a>.</p>

Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to [RabMAb<sup>®</sup> patents](#).

## 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C. Do Not Freeze.
存储溶液	pH: 7.20 Constituent: PBS
无载体	是
纯度	Protein A purified
克隆	单克隆
克隆编号	Y299
同种型	IgG

## 应用

**The Abpromise guarantee**      **Abpromise<sup>™</sup>承诺保证使用ab219724于以下的经测试应用**

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC/IF		Use at an assay dependent concentration.
Flow Cyt (Intra)		Use at an assay dependent concentration. <b>ab199376</b> - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
IP		Use at an assay dependent concentration.
WB		Use at an assay dependent concentration. Detects a band of approximately 12 kDa (predicted molecular weight: 12 kDa).
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
ChIP		Use at an assay dependent concentration.

## 靶标

**功能**      Ubiquitin-like protein that can be covalently attached to proteins as a monomer or a lysine-linked polymer. Covalent attachment via an isopeptide bond to its substrates requires prior activation by the E1 complex SAE1-SAE2 and linkage to the E2 enzyme UBE2I, and can be promoted by E3 ligases such as PIAS1-4, RANBP2 or CBX4. This post-translational modification on lysine residues of proteins plays a crucial role in a number of cellular processes such as nuclear

transport, DNA replication and repair, mitosis and signal transduction. Involved for instance in targeting RANGAP1 to the nuclear pore complex protein RANBP2. Polymeric SUMO1 chains are also susceptible to polyubiquitination which functions as a signal for proteasomal degradation of modified proteins. May also regulate a network of genes involved in palate development.

#### 疾病相关

Defects in SUMO1 are the cause of non-syndromic orofacial cleft type 10 (OFC10) [MIM:613705]; also called non-syndromic cleft lip with or without cleft palate 10. OFC10 is a birth defect consisting of cleft lips with or without cleft palate. Cleft lips are associated with cleft palate in two-third of cases. A cleft lip can occur on one or both sides and range in severity from a simple notch in the upper lip to a complete opening in the lip extending into the floor of the nostril and involving the upper gum. Note=A chromosomal aberration involving SUMO1 is the cause of OFC10. Translocation t(2;8)(q33.1;q24.3). The breakpoint occurred in the SUMO1 gene and resulted in haploinsufficiency confirmed by protein assays.

#### 序列相似性

Belongs to the ubiquitin family. SUMO subfamily.  
Contains 1 ubiquitin-like domain.


#### 翻译后修饰

Cleavage of precursor form by SENP1 or SENP2 is necessary for function.  
Polymeric SUMO1 chains undergo polyubiquitination by RNF4.

#### 细胞定位

Nucleus membrane. Nucleus speckle. Cytoplasm. Recruited by BCL11A into the nuclear body.

#### 图片




**ab32058** immunoprecipitating Sumo 1. 10µg of NIH/3T3 (Mouse embryonic fibroblast) cell lysate was incubated with primary antibody at a dilution of 1/20 and VeriBlot for IP Detection Reagent (HRP) (**ab131366**) at a dilution of 1/1000.

Lane 1: NIH/3T3 whole cell lysate 10ug

Lane 2: **ab32058** IP in NIH/3T3 whole cell lysate

Lane 3: Rabbit monoclonal IgG (**ab172730**) instead of **ab32058** in NIH/3T3 (Mouse embryonic fibroblast) whole cell lysate


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32058**).



Chromatin was prepared from SK-OV-3 (Human ovarian cancer cell line) cells according to the Abcam X-ChIP protocol. Cells were fixed with formaldehyde for 10 minutes. The ChIP was performed with 25µg of chromatin, 5µg of **ab32058** (blue), and 20µl of Anti rabbit IgG sepharose beads. 5µg of rabbit normal IgG was added to the beads control (yellow). The immunoprecipitated DNA was quantified by real time PCR (Sybr green approach).

ChIP was performed according to the literature (PMID: 23770046).


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32058**).



**ab32058** staining Sumo 1 in rat stomach tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with paraformaldehyde and antigen retrieval was by heat mediation in a EDTA buffer. Samples were incubated with primary antibody at a dilution of 1/250. A goat anti-rabbit IgG H&L (HRP) **ab97051** was used as the secondary antibody at a dilution of 1/500.

Negative control 1: PBS in place of primary antibody.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32058**).




Immunocytochemistry/ Immunofluorescence - Anti-Sumo 1 antibody [Y299] - BSA and Azide free (ab219724)

Clone Y299 (ab219724) has been successfully conjugated by Abcam. This image was generated using Anti-Sumo 1 antibody [Y299] (Alexa Fluor® 488). Please refer to [ab196310](#) for protocol details.

[ab196310](#) staining Sumo 1 in NIH3T3 cells. The cells were fixed with 4% formaldehyde (10 min), permeabilised in 0.1% Triton X-100 for 5 minutes and then blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1%PBS-Tween for 1h. The cells were then incubated with [ab196310](#) at 1/50 dilution (shown in green) and [ab195889](#), Mouse monoclonal [DM1A] to alpha Tubulin (Alexa Fluor® 594, shown in red) at 1/167 dilution overnight at +4°C. Nuclear DNA was labelled in blue with DAPI.

This product gave a positive signal in 100% methanol (5 min) fixed NIH3T3 cells under the same testing conditions.

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).




Immunocytochemistry/ Immunofluorescence - Anti-Sumo 1 antibody [Y299] - BSA and Azide free (ab219724)

Clone Y299 (ab219724) has been successfully conjugated by Abcam. This image was generated using Anti-Sumo 1 antibody [Y299] (Alexa Fluor® 647). Please refer to [ab196533](#) for protocol details.

[ab196533](#) staining Sumo 1 in NIH3T3 cells. The cells were fixed with 4% formaldehyde (10 min), permeabilised in 0.1% Triton X-100 for 5 minutes and then blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1%PBS-Tween for 1h. The cells were then incubated with [ab196533](#) at 1/50 dilution (shown in red) and [ab195887](#), Mouse monoclonal [DM1A] to alpha Tubulin (Alexa Fluor® 594, shown in green) at 1/167 dilution overnight at +4°C. Nuclear DNA was labelled in blue with DAPI.

This product gave a positive signal in 100% methanol (5 min) fixed NIH3T3 cells under the same testing conditions.


Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).



Immunocytochemistry/ Immunofluorescence - Anti-Sumo 1 antibody [Y299] - BSA and Azide free (ab219724)

**ab32058** staining Sumo 1 in HeLa (Human epithelial cell line from cervix adenocarcinoma) cells by ICC/IF (Immunocytochemistry/immunofluorescence). Cells were fixed with 4% Paraformaldehyde and permeabilized with 0.1% Triton X-100. Samples were incubated with primary antibody at a dilution of 1/500. A goat anti rabbit IgG (Alexa Fluor® 488) (**ab150077**) was used as the secondary antibody at a dilution of 1/1000. **ab195889** was used as a counterstain for primary antibody **ab133645** at 1/200. DAPI was used as a nuclear counterstain and PBS as a negative control.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32058**).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Sumo 1 antibody [Y299] - BSA and Azide free (ab219724)

**ab32058** staining Sumo 1 in mouse kidney tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with paraformaldehyde and antigen retrieval was by heat mediation in a EDTA buffer. Samples were incubated with primary antibody at a dilution of 1/250. A goat anti-rabbit IgG H&L (HRP) **ab97051** was used as the secondary antibody at a dilution of 1/500.

**Negative control 1:** PBS in place of primary antibody.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32058**).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Sumo 1 antibody [Y299] - BSA and Azide free (ab219724)

**ab32058** staining Sumo 1 in human bladder carcinoma tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with paraformaldehyde and antigen retrieval was by heat mediation in a EDTA buffer. Samples were incubated with primary antibody at a dilution of 1/250. A goat anti-rabbit IgG H&L (HRP) **ab97051** was used as the secondary antibody at a dilution of 1/500.

Negative control 1: PBS in place of primary antibody.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32058**).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Sumo 1 antibody [Y299] - BSA and Azide free (ab219724)

**ab32058** staining Sumo 1 in human endometrium tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with paraformaldehyde and antigen retrieval was by heat mediation in a EDTA buffer. Samples were incubated with primary antibody at a dilution of 1/250. A goat anti-rabbit IgG H&L (HRP) **ab97051** was used as the secondary antibody at a dilution of 1/500.

**Negative control 1:** PBS in place of primary antibody.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32058**).




Flow Cytometry (Intracellular) - Anti-Sumo 1 antibody [Y299] - BSA and Azide free (ab219724)

**ab32058** staining Sumo 1 in HeLa (Human epithelial cell line from cervix adenocarcinoma) cells by intracellular flow cytometry. Cells were fixed with 4% paraformaldehyde and the sample was incubated with the primary antibody at a dilution of 1/20. A goat anti rabbit IgG (Alexa Fluor<sup>®</sup> 488) at a dilution of 1/2000 was used as the secondary antibody.

Isotype control: Rabbit monoclonal IgG (Black)

Unlabelled control: Cell without incubation with primary antibody and secondary antibody (Blue)

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32058**).




Flow Cytometry (Intracellular) - Anti-Sumo 1 antibody [Y299] - BSA and Azide free (ab219724)

Intracellular Flow Cytometry analysis of HeLa (human cervix adenocarcinoma) cells labelling Sumo 1 with **ab32058** at 1/20 (red). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. An Alexa Fluor<sup>®</sup>488-conjugated goat anti-rabbit IgG (1/2000) was used as the secondary antibody. Black - Isotype control, rabbit monoclonal IgG. Blue - Unlabelled control, cells without incubation with primary and secondary antibodies.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32058**).






Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Sumo 1 antibody [Y299] - BSA and Azide free (ab219724)

IHC of paraffin-embedded human lung carcinoma using anti-SUMO 1 ([ab32058](#)) diluted 1:250


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab32058](#)).



Immunocytochemistry/ Immunofluorescence - Anti-Sumo 1 antibody [Y299] - BSA and Azide free (ab219724)

Immunofluorescent staining of HeLa (Human epithelial cell line from cervix adenocarcinoma) cells using anti-SUMO 1 ([ab32058](#)) diluted 1/250.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab32058](#)).



Immunocytochemistry/ Immunofluorescence - Anti-Sumo 1 antibody [Y299] - BSA and Azide free (ab219724)

Image from Cuchet-Lourenço D et al. PLoS Pathog. 2011 Jul;7(7):e1002123. Epub 2011 Jul 14. Fig 9; doi:10.1371/journal.ppat.1002123; July 14 2011 PLoS Pathog 7(7): e1002123.

Immunofluorescence analysis of ICP0-null mutant HSV-1 infected HepaRG cells, staining Sumo1 (green) with [ab32058](#). An AlexaFluor®-conjugated goat anti-rabbit IgG was used as the secondary antibody.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab32058](#)).

### Why choose a recombinant antibody?



**Research with confidence**  
Consistent and reproducible results



**Long-term and scalable supply**  
Recombinant technology



**Success from the first experiment**  
Confirmed specificity



**Ethical standards compliant**  
Animal-free production

Anti-Sumo 1 antibody [Y299] - BSA and Azide free  
(ab219724)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors