


Anti-SUZ12 antibody [EPR5234(N)] - CHIP Grade ab175187

敲除验证
重组
RabMAb

★★★★★
[1 Abreviews](#)
[5 References](#)
[12 图像](#)

概述

产品名称	Anti-SUZ12抗体[EPR5234(N)] - CHIP Grade
描述	兔单克隆抗体[EPR5234(N)] to SUZ12 - CHIP Grade
宿主	Rabbit
经测试应用	适用于: Flow Cyt (Intra), ChIP, ChIC/CUT&RUN-seq, WB, ICC/IF, IP 不适用于: IHC-P
种属反应性	与反应: Mouse, Human 预测可用于: Rat 
免疫原	Synthetic peptide within Human SUZ12 aa 50-150 (Cysteine residue). The exact sequence is proprietary. Database link: Q15022
阳性对照	WB: HAP1, Caco2, MCF7, SW480 and 293T cell lysate. IP: HeLa whole cell lysate. ChIP: HeLa and F9 cells. ICC/IF: MCF7 cells. Flow Cyt (intra): HeLa cells. ChIC/CUT&RUN-Seq: HeLa cells.
常规说明	This product is a recombinant monoclonal antibody, which offers several advantages including: <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production For more information see here . Our RabMAb [®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents .

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
存储溶液	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 9% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA, 50% Tissue culture supernatant

纯度	Protein A purified
克隆	单克隆
克隆编号	EPR5234(N)
同种型	IgG

应用

The Abpromise guarantee **Abpromise™** 承诺保证使用 ab175187 于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度；实际最佳的稀释度/浓度应由使用者检定。


应用	Ab评论	说明
Flow Cyt (Intra)		1/2000.
ChIP	★☆☆☆☆ (1)	Use at an assay dependent concentration.
ChIC/CUT&RUN-seq		Use at an assay dependent concentration. 5µg
WB		1/1000 - 1/10000. Predicted molecular weight: 83 kDa.
ICC/IF		1/50 - 1/100.
IP		1/10 - 1/100.

应用说明 Is unsuitable for IHC-P.

靶标

功能	Polycomb group (PcG) protein. Component of the PRC2/EED-EZH2 complex, which methylates 'Lys-9' (H3K9me) and 'Lys-27' (H3K27me) of histone H3, leading to transcriptional repression of the affected target gene. The PRC2/EED-EZH2 complex may also serve as a recruiting platform for DNA methyltransferases, thereby linking two epigenetic repression systems. Genes repressed by the PRC2/EED-EZH2 complex include HOXC8, HOXA9, MYT1 and CDKN2A.
组织特异性	Overexpressed in breast and colon cancer.
疾病相关	Note=A chromosomal aberration involving SUZ12 may be a cause of endometrial stromal tumors. Translocation t(7;17)(p15;q21) with JAZF1. The translocation generates the JAZF1-SUZ12 oncogene consisting of the N-terminus part of JAZF1 and the C-terminus part of SUZ12. It is frequently found in all cases of endometrial stromal tumors, except in endometrial stromal sarcomas, where it is rarer.
序列相似性	Belongs to the VEFS (VRN2-EMF2-FIS2-SU(Z)12) family. Contains 1 C2H2-type zinc finger.
发展阶段	Expressed at low levels in quiescent cells. Expression rises at the G1/S phase transition.
细胞定位	Nucleus.


图片



ChIC/CUT&RUN was performed using a pAG-MNase at a final concentration of 700 ng/mL, 2×10^5 HeLa (Human cervix adenocarcinoma epithelial cell line) cells and 5µg of ab175187 [EPR5234(N)]. The resulting DNA was sequenced on the Illumina NovaSeq 6000 to a depth of 10 million reads. The negative IgG control **ab172730** is also shown.

Additional screenshots of mapped reads can be downloaded [here](#).

The University of Geneva owns patents relevant to ChIC (Chromatin Immuno-Cleavage) methods.



All lanes : Anti-SUZ12 antibody [EPR5234(N)] - ChIP Grade (ab175187) at 1/1000 dilution

Lane 1 : Wild-type HAP1 cell lysate

Lane 2 : SUZ12 knockout HAP1 cell lysate

Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.

Predicted band size: 83 kDa

Observed band size: 90 kDa

Lanes 1 - 2: Merged signal (red and green). Green - ab175187 observed at 90 kDa. Red - loading control **ab7291** (Mouse anti-Alpha Tubulin [DM1A]) observed at 55 kDa.

ab175187 was shown to react with SUZ12 in wild-type HAP1 cells in Western blot with loss of signal observed in SUZ12 knockout sample. Wild-type HAP1 and SUZ12 knockout cell lysates were subjected to SDS-PAGE. Membranes were blocked in 3 % milk in TBS-T (0.1 % Tween®) before incubation with ab175187 and **ab7291** (Mouse anti-Alpha Tubulin [DM1A]) overnight at 4 °C at a 1 in 1000 dilution and a 1 in 20000 dilution respectively. Blots were incubated with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 h at room temperature before imaging.



Immunoprecipitation - Anti-SUZ12 antibody
[EPR5234(N)] - ChIP Grade (ab175187)

ab175187 (purified) at 1/20 dilution (16 µg/mL)
immunoprecipitating SUZ12 in HeLa (human cervix adenocarcinoma epithelial cell) whole cell lysate 10 µg.
Lane 1 (input): HeLa (human cervix adenocarcinoma epithelial cell) whole cell lysate 10 µg
Lane 2 (+): ab175187 & HeLa whole cell lysate
Lane 3 (-): Rabbit monoclonal IgG (**ab172730**) instead of ab175187 in HeLa whole cell lysate
For western blotting, ab175187 at 1/500 dilution (0.636 µg/mL) and veriBlot for IP secondary antibody (HRP) (**ab131366**) at 1/1000 dilution was used.

Blocking and diluting buffer: 5% NFDM /TBST.



ChIP - Anti-SUZ12 antibody [EPR5234(N)] - ChIP Grade (ab175187)


Chromatin was prepared from HeLa cells according to the Abcam Dual X-ChIP protocol*. Cells were fixed with EGS for 30 minutes, then formaldehyde for 10 minutes.

The ChIP was performed with 25 µg of chromatin, 5 µg of ab175187 (red), and 20 µl of Protein A/G sepharose beads. 5 µg of rabbit normal IgG was added to the beads control (gray). The immunoprecipitated DNA was quantified by real time PCR (Sybr green approach).

Primers and probes are located in the first kb of the transcribed region.

*[http://www.abcam.com/resources?](http://www.abcam.com/resources?keywords=X%20ChIP%20protocol)

keywords=X%20ChIP%20protocol



Western blot - Anti-SUZ12 antibody [EPR5234(N)] - ChIP Grade (ab175187)

Lane 1: Wild-type HAP1 cell lysate (20 µg)


Lane 2: SUZ12 knockout HAP1 cell lysate (20 µg)

Lane 3: Caco2 cell lysate (20 µg)

Lane 4: MCF7 cell lysate (20 µg)


Lanes 1 - 4: Merged signal (red and green). Green - ab175187 observed at 100 kDa. Red - loading control, **ab8245**, observed at 37 kDa.

ab175187 was shown to specifically react with SUZ12 in wild-type HAP1 cells along with additional cross-reactive bands. No band was observed when SUZ12 knockout samples were used. Wild-type and SUZ12 knockout samples were subjected to SDS-PAGE. ab175187 and **ab8245** (loading control to GAPDH) were both 1/1000 and 1/10,000 respectively and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1/10,000 dilution for 1hr at room temperature before imaging.




Immunocytochemistry/ Immunofluorescence - Anti-SUZ12 antibody [EPR5234(N)] - ChIP Grade (ab175187)

Immunofluorescence analysis of MCF-7 cells labeling SUZ12 with ab175187 at a 1/50 dilution.



Flow Cytometry (Intracellular) - Anti-SUZ12 antibody [EPR5234(N)] - ChIP Grade (ab175187)

Intracellular Flow Cytometry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling SUZ12 (red) with purified ab175187 at a 1/2000 dilution (1ug/mL). Cells were fixed with 4% paraformaldehyde and permeabilized with 90% methanol. A goat anti rabbit IgG (Alexa Fluor® 488) (**ab150077**) was used as the secondary antibody at a 1/2000 dilution. Black - Rabbit monoclonal IgG (Black) (**ab172730**). Blue (unlabeled control) - Cell without incubation with primary antibody and secondary antibody (Blue).



Western blot - Anti-SUZ12 antibody [EPR5234(N)] - ChIP Grade (ab175187)

All lanes : Anti-SUZ12 antibody [EPR5234(N)] - ChIP Grade (ab175187) at 1/1000 dilution

Lane 1 : Wild-type HeLa cell lysate

Lane 2 : SUZ12 CRISPR/Cas9 edited HeLa cell lysate

Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.


Predicted band size: 83 kDa

Observed band size: 100 kDa

Lanes 1-2: Merged signal (red and green). Green - ab175187 observed at 100 kDa. Red - Anti-GAPDH antibody [6C5] - Loading Control (**ab8245**) observed at 37 kDa.

ab175187 was shown to react with SUZ12 in wild-type HeLa cells in western blot. The band observed in CRISPR/Cas9 edited cell line **ab264983** (CRISPR/Cas9 edited cell lysate **ab257721**) lane below 100kDa may represent truncated forms and cleaved fragments. This has not been investigated further. Wild-type HeLa and SUZ12 CRISPR/Cas9 edited HeLa cell lysates were subjected to SDS-PAGE. Membrane was blocked for 1 hour at room temperature in 0.1% TBST with 3% non-fat dried milk. ab175187 and Anti-GAPDH antibody [6C5] - Loading Control (**ab8245**) were incubated overnight at 4°C at a 1 in 1000 dilution and a 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L

(IRDye[®]800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye[®]680RD) preadsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Western blot - Anti-SUZ12 antibody [EPR5234(N)] - ChIP Grade (ab175187)

All lanes : Anti-SUZ12 antibody [EPR5234(N)] - ChIP Grade (ab175187) at 1/1000 dilution

Lane 1 : SW480 cell lysates

Lane 2 : HeLa cell lysates

Lane 3 : MCF-7 cell lysates

Lane 4 : 293T cell lysates


Lysates/proteins at 10 µg per lane.

Predicted band size: 83 kDa



ChIP - Anti-SUZ12 antibody [EPR5234(N)] - ChIP Grade (ab175187)

Chromatin was prepared from F9 cells according to the Abcam X-ChIP protocol. Cells were fixed with formaldehyde for 10 minutes. The ChIP was performed with 25 µg of chromatin, 5 µg of ab175187 (red), and 20 µl of Protein A/G sepharose beads. 5 µg of rabbit normal IgG was added to the beads control (gray). The immunoprecipitated DNA was quantified by real time PCR (Taqman approach for active and inactive loci). Primers and probes are located in the first kb of the transcribed region.



Western blot analysis on immunoprecipitation pellet from HeLa cell lysate using ab175187 at a 1/10 dilution.

Immunoprecipitation - Anti-SUZ12 antibody
[EPR5234(N)] - ChIP Grade (ab175187)

Why choose a recombinant antibody?

 Research with confidence Consistent and reproducible results	 Long-term and scalable supply Recombinant technology
 Success from the first experiment Confirmed specificity	 Ethical standards compliant Animal-free production

Anti-SUZ12 antibody [EPR5234(N)] - ChIP Grade
(ab175187)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Promise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors