

Anti-Syntenin antibody [EPR8102] ab133267

敲除验证 重组 RabMAb


★★★★★ 1 Abreviews 68 References 14 图像

概述

产品名称	Anti-Syntenin抗体[EPR8102]
描述	兔单克隆抗体[EPR8102] to Syntenin
宿主	Rabbit
经测试应用	适用于: Flow Cyt (Intra), WB, IP, IHC-P, ICC/IF
种属反应性	与反应: Human
免疫原	Synthetic peptide within Human Syntenin aa 1-100. The exact sequence is proprietary. Database link: <a href="#">O00560</a>
阳性对照	Human fetal brain lysate, Human fetal heart lysate, Human placenta lysate, HeLa, 293T and A549 (Human lung carcinoma cell line) cell lysates; Human brain tissue
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"><li>- High batch-to-batch consistency and reproducibility</li><li>- Improved sensitivity and specificity</li><li>- Long-term security of supply</li><li>- Animal-free production</li></ul> <p>For more information <a href="#">see here</a>.</p> <p>Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a>.</p> <p>Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
解离常数 (K <sub>D</sub> )	K <sub>D</sub> = 1.24 x 10 <sup>-10</sup> M



[Learn more about Kp](#)

存储溶液	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol, 0.05% BSA
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR8102
同种型	IgG

## 应用

**The Abpromise guarantee**      **Abpromise™** 承诺保证使用 ab133267 于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab 评论	说明
Flow Cyt (Intra)		1/50. <b>ab172730</b> - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody. For unpurified use at 1/1000 - 1/10000.
WB	★★★★★ (1)	1/1000 - 1/10000. Detects a band of approximately 32 kDa (predicted molecular weight: 32 kDa).
IP		1/10 - 1/100.
IHC-P		1/50 - 1/100. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. See <b><u>IHC antigen retrieval protocols</u></b> .
ICC/IF		1/50 - 1/200.

## 靶标

功能	Seems to function as an adapter protein. In adherens junctions may function to couple syndecans to cytoskeletal proteins or signaling components. Seems to couple transcription factor SOX4 to the IL-5 receptor (IL5RA). May also play a role in vesicular trafficking. Seems to be required for the targeting of TGFA to the cell surface in the early secretory pathway.
组织特异性	Widely expressed. Expressed in fetal kidney, liver, lung and brain. In adult highest expression in heart and placenta.
序列相似性	Contains 2 PDZ (DHR) domains.
翻译后修饰	Phosphorylated on tyrosine residues.
细胞定位	Cell junction > focal adhesion. Cell junction > adherens junction. Cell membrane. Endoplasmic reticulum membrane. Nucleus. Melanosome. Cytoplasm > cytosol. Cytoplasm > cytoskeleton. Mainly membrane-associated. Localized to adherens junctions, focal adhesions and endoplasmic reticulum. Colocalized with actin stress fibers. Also found in the nucleus. Identified by mass

## 图片



Western blot - Anti-Syntenin antibody [EPR8102]  
(ab133267)

**Lane 1:** Wild-type HAP1 whole cell lysate (20 µg)


**Lane 2:** Syntenin knockout HAP1 whole cell lysate (20 µg)

**Lane 3:** A549 whole cell lysate (20 µg)

**Lane 4:** HeLa whole cell lysate (20 µg)


**Lanes 1 - 4:** Merged signal (red and green). Green - ab133267 observed at 32 kDa. Red - loading control, **ab130007**, observed at 130 kDa.

ab133267 was shown to recognize Syntenin in wild-type HAP1 cells as signal was lost at the expected MW in Syntenin knockout cells. Additional cross-reactive bands were observed in the wild-type and knockout cells. Wild-type and Syntenin knockout samples were subjected to SDS-PAGE. ab133267 and **ab130007** (Mouse anti-Vinculin loading control) were incubated overnight at 4°C at 1/1000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed **ab216773** and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed **ab216776** secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Syntenin antibody [EPR8102] (ab133267)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human hepatocellular carcinoma tissue labeling Syntenin with purified ab133267 at 1/50 dilution. Heat mediated antigen retrieval was performed using EDTA buffer pH 9. **ab97051**, a HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500). Negative control using PBS instead of primary antibody. Counterstained with hematoxylin



Western blot - Anti-Syntenin antibody [EPR8102] (ab133267)

Anti-Syntenin antibody [EPR8102] (ab133267) at 1/2000 dilution (purified) + A549 (Human lung carcinoma cell line) whole cell lysate at 15 µg


### Secondary

Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution (Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated)

**Predicted band size:** 32 kDa


**Observed band size:** 32 kDa

Blocking and dilution buffer: 5% NFDM/TBST



Flow Cytometry (Intracellular) - Anti-Syntenin antibody [EPR8102] (ab133267)

Intracellular Flow Cytometry analysis of HeLa (Human epithelial cell line from cervix adenocarcinoma) labeling Syntenin with purified ab133267 at 1/50 (red). Cells were fixed with 4% paraformaldehyde. A goat anti rabbit IgG (Alexa Fluor® 488) 1/2000 was used as the secondary antibody. Black - Isotype control, rabbit monoclonal IgG. Blue - Unlabelled control, cells without incubation with primary and secondary antibodies.



Immunoprecipitation - Anti-Syntenin antibody  
[EPR8102] (ab133267)

ab133267 (purified) at 1/40 immunoprecipitating Syntenin in HeLa whole cell lysate.


Lane 1 (input): HeLa whole cell lysate (10µg)

Lane 2 (+): ab133267 + HeLa whole cell lysate.

Lane 3 (-): Rabbit monoclonal IgG (**ab172730**) instead of ab133267 in HeLa whole cell lysate.


Blocking buffer and concentration: 5% NFDM/TBST.

Diluting buffer and concentration: 5% NFDM /TBST.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Syntenin antibody  
[EPR8102] (ab133267)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human cerebral cortex tissue labeling Syntenin with purified ab133267 at 1/50 dilution. Heat mediated antigen retrieval was performed using EDTA buffer pH 9. **ab97051**, a HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500). Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.



**All lanes :** Anti-Syntenin antibody [EPR8102] (ab133267) at 1/2000 dilution (purified)

**Lane 1 :** Human placenta lysate

**Lane 2 :** Human fetal brain lysate

Lysates/proteins at 15 µg per lane.


#### Secondary

**All lanes :** Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG at 1/2000 dilution

**Predicted band size:** 32 kDa

**Observed band size:** 32 kDa

Blocking and dilution buffer: 5% NFDM/TBST



Anti-Syntenin antibody [EPR8102] (ab133267) at 1/10000 dilution (purified) + HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate at 20 µg


#### Secondary

Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG at 1/2000 dilution

**Predicted band size:** 32 kDa

**Observed band size:** 32 kDa


Blocking and dilution buffer: 5% NFDM/TBST



Immunocytochemistry/ Immunofluorescence - Anti-Syntenin antibody [EPR8102] (ab133267)

ab133267 staining Syntenin in HeLa (human cervix adenocarcinoma) cells by ICC/IF (Immunocytochemistry/immunofluorescence). Cells were fixed with 4% Paraformaldehyde and permeabilized with 0.1% Triton X-100. Samples were incubated with primary antibody at a dilution of 1/500. A goat anti rabbit IgG (Alexa Fluor® 488) ([ab150077](#)) was used as the secondary antibody at a dilution of 1/1000. DAPI was used as a nuclear counterstain.


**Negative control 1:** PBS only.



Flow Cytometry (Intracellular) - Anti-Syntenin antibody [EPR8102] (ab133267)

Overlay histogram showing SHSY-5Y cells stained with unpurified ab133267 (red line). The cells were fixed with 4% paraformaldehyde (10 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab133267, 1/10000 dilution) for 30 min at 22°C. The secondary antibody used was Alexa Fluor® 488 goat anti-rabbit IgG (H&L) ([ab150077](#)) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (0.1µg/1x10<sup>6</sup> cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter. This antibody gave a positive signal in SHSY-5Y cells fixed with 80% methanol (5 min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.





**All lanes :** Anti-Syntenin antibody [EPR8102] (ab133267) at 1/1000 dilution (unpurified)

**Lane 1 :** Human fetal brain lysate

**Lane 2 :** 293T cell lysate

**Lane 3 :** Human fetal heart lysate

**Lane 4 :** HeLa cell lysate


Lysates/proteins at 10 µg per lane.

### Secondary

**All lanes :** Goat anti-rabbit HRP conjugated antibody at 1/2000 dilution

**Predicted band size:** 32 kDa

**Observed band size:** 32 kDa




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Syntenin antibody [EPR8102] (ab133267)

Immunohistochemical analysis of Syntenin in paraffin embedded Human brain tissue, using unpurified ab133267 at a 1/50 dilution.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.





BI-RD Scanning - Anti-Syntenin antibody [EPR8102]  
(ab133267)

Equilibrium disassociation constant ( $K_D$ )

Learn more about  $K_D$

[Click here to learn more about  \$K\_D\$](#)

### Why choose a recombinant antibody?



**Research with confidence**  
Consistent and reproducible results



**Long-term and scalable supply**  
Recombinant technology



**Success from the first experiment**  
Confirmed specificity



**Ethical standards compliant**  
Animal-free production

Anti-Syntenin antibody [EPR8102] (ab133267)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

#### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors