

Anti-TAGLN/Transgelin antibody ab14106

★★★★★ [27 Abreviews](#) [383 References](#) [6 图像](#)

概述

产品名称	Anti-TAGLN/Transgelin抗体
描述	兔多克隆抗体to TAGLN/Transgelin
宿主	Rabbit
特异性	The immunogen used to generate this TAGLN/Transgelin antibody (ab14106) shares 75% homology with family members TAGLN2 and TAGLN3. Cross-reactivity with these proteins has not been confirmed experimentally.
经测试应用	适用于: ICC/IF, WB
种属反应性	与反应: Mouse, Rat, Human
免疫原	Synthetic peptide within Mouse TAGLN/Transgelin aa 150 to the C-terminus conjugated to keyhole limpet haemocyanin. The exact sequence is proprietary. (Peptide available as ab16067)
阳性对照	WB: Human primary smooth muscle cells; HeLa whole cell lysate; Mouse colon tissue lysate; Rat colon tissue lysate. ICC/IF: HeLa cells; mouse muscle cells.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	pH: 7.40 Preservative: 0.02% Sodium azide Constituent: PBS
	Batches of this product that have a concentration < 1mg/ml may have BSA added as a stabilising agent. If you would like information about the formulation of a specific lot, please contact our

	scientific support team who will be happy to help.
纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

应用

The Abpromise guarantee Abpromise™ 承诺保证使用 ab14106 于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab 评论	说明
ICC/IF	★★★★★ (8)	Use a concentration of 1 - 5 µg/ml.
WB	★★★★★ (12)	Use a concentration of 1 µg/ml. Detects a band of approximately 23 kDa (predicted molecular weight: 23 kDa).

靶标

功能	Actin cross-linking/gelling protein (By similarity). Involved in calcium interactions and contractile properties of the cell that may contribute to replicative senescence.
序列相似性	Belongs to the calponin family. Contains 1 calponin-like repeat. Contains 1 CH (calponin-homology) domain.
细胞定位	Cytoplasm.

图片



All lanes : Anti-TAGLN/Transgelin antibody (ab14106) at 1/1000 dilution

All lanes : Lysates prepared from human primary smooth muscle cells

Lysates/proteins at 30 µg per lane.

Secondary

All lanes : HRP-conjugated bovine polyclonal to rabbit IgG at 1/2000 dilution


Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 23 kDa

Observed band size: 23 kDa

Exposure time: 2 minutes




Immunocytochemistry/ Immunofluorescence - Anti-TAGLN/Transgelin antibody (ab14106)

ab14106 stained HeLa (Human epithelial cell line from cervix adenocarcinoma) cells.

The cells were fixed in 100% methanol for 5 minutes, permeabilized in 0.1% PBS-Triton X-100 for 5 minutes and then incubated in 1% BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1 hour at room temperature to block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab14106 at 1 µg/ml) overnight at +4°C. The secondary antibody (pseudo-colored green) was **ab150081** Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed used at a 1/1000 dilution for 1 hour at room temperature. Alexa Fluor® 594 WGA was used to label plasma membranes (pseudo-colored red) at a 1/200 dilution for 1 hour at room temperature.

DAPI was used to stain the cell nuclei (pseudo-colored blue) at a concentration of 1.43µM for 1 hour at room temperature.




Immunocytochemistry/ Immunofluorescence - Anti-TAGLN/Transgelin antibody (ab14106)

This image is courtesy of an anonymous Abreview.

ab14106 staining SM22 alpha in mouse muscle cells by ICC/IF (Immunocytochemistry/immunofluorescence).

Cells were fixed with paraformaldehyde, permeabilized with 0.1% Triton X-100 and blocked with 10% serum for 30 minutes at 25°C. Samples were incubated with primary antibody (1/100 in PBS) for 18 hours at 4°C. An Alexa Fluor® 488-conjugated goat anti-rabbit IgG polyclonal was used as the secondary antibody at a dilution of 1/1000.



All lanes : Anti-TAGLN/Transgelin antibody (ab14106) at 1 µg/ml

Lane 1 : HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate

Lane 2 : HeLa nuclear extract

Lane 3 : Human skeletal muscle tissue lysate

Lysates/proteins at 20 µg per lane.


Secondary

All lanes : Alexa Fluor Goat polyclonal to Rabbit IgG at 1/10000 dilution

Predicted band size: 23 kDa

Observed band size: 23 kDa

The band was completely abolished by peptide blocking with **ab16067** (immunizing peptide) - not shown.



Western blot - Anti-TAGLN/Transgelin antibody
(ab14106)

All lanes : Anti-TAGLN/Transgelin antibody (ab14106) at 1 µg/ml

Lane 1 : Human colon tissue lysate - total protein ([ab30051](#))

Lane 2 : Mouse colon tissue lysate

Lane 3 : Rat colon tissue lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) preadsorbed at
1/50000 dilution

Developed using the ECL technique.

Performed under reducing conditions.


Predicted band size: 23 kDa

Observed band size: 23 kDa

Additional bands at: 115 kDa, 20 kDa, 36 kDa. We are unsure as
to the identity of these extra bands.

Exposure time: 5 seconds

This blot was produced using a 4-12% Bis-tris gel under the MES buffer system. The gel was run at 200V for 35 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour using 2% Bovine Serum Albumin before being incubated with ab14106 overnight at 4°C. Antibody binding was detected using an anti-rabbit antibody conjugated to HRP, and visualised using ECL development solution [ab133406](#).



Anti-TAGLN/Transgelin antibody (ab14106) at 1 µg/ml +
Recombinant Human TAGLN/Transgelin protein (**ab101469**) at
0.01 µg

Secondary

Goat Anti-Rabbit IgG H&L (HRP) preadsorbed (**ab97080**) at
1/5000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 23 kDa

Exposure time: 30 seconds

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors