# abcam

### **Product datasheet**

## Anti-TBLR1/TBL1XR1 antibody [4F3-A8-D9] ab117761

<u>2 References</u> 7 图像

概述		
产品名称	Anti-TBLR1/TBL1XR1 <b>抗体</b> [4F3-A8-D9]	
描述	<b>小鼠</b> 单 <b>克隆抗体</b> [4F3-A8-D9] to TBLR1/TBL1XR1	
宿主	Mouse	
经 <b>测</b> 试应 <b>用</b>	适用于: WB, IHC-P, ICC/IF	
种属反应性	<b>与反应:</b> Mouse, Human, African green monkey	
免疫原	Recombinant fragment corresponding to Human TBLR1/TBL1XR1. Database link: <b>Q9BZK7</b>	
<b>阳性</b> 对 <b>照</b>	WB: K562 whole cell lysate, Mouse heart lysate, Mouse brain lysate ICC: HeLa cells IHC-P: Human breast cancer	
<b>常</b> 规说 <b>明</b>	This product was changed from ascites to tissue culture supernatant on 17.08.2018. Please note that the dilutions may need to be adjusted accordingly. If you have any questions, please do not hesitate to contact our scientific support team.	
	The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.	
	If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As	

性能	
形式	Liquid
存 <b>放</b> 说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid repeated freeze / thaw cycles.
存储溶液	pH: 7.40 Preservative: 0.03% Proclin 300 Constituent: 50% Glycerol
纯 <b>度</b>	Concentrated Culture Supernatant
克隆	单 <b>克隆</b>
<b>克隆</b> 编号	4F3-A8-D9

#### 应用


#### The Abpromise guarantee Abpromise ™承诺保证使用ab117761于以下的经测试应用

#### "应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。


应用	Ab评论	说明
WB		1/1000. Detects a band of approximately 60 kDa (predicted molecular weight: 56 kDa).
IHC-P		1/500 - 1/1000.
ICC/IF		1/150 - 1/200.

<b>靶</b> 标	
功能	F-box-like protein involved in the recruitment of the ubiquitin/19S proteasome complex to nuclear receptor-regulated transcription units. Plays an essential role in transcription activation mediated by nuclear receptors. Probably acts as integral component of the N-Cor corepressor complex that mediates the recruitment of the 19S proteasome complex, leading to the subsequent proteasomal degradation of N-Cor complex, thereby allowing cofactor exchange, and transcription activation.
组织 <b>特异性</b>	Ubiquitous.
序列相似性	Belongs to the WD repeat EBI family. Contains 1 F-box-like domain. Contains 1 LisH domain. Contains 8 WD repeats.
结 <b>构域</b>	The F-box-like domain is related to the F-box domain, and apparently displays the same function as component of ubiquitin E3 ligase complexes.
<b>翻</b> 译后 <b>修</b> 饰	Phosphorylated upon DNA damage, probably by ATM or ATR.
细胞定位	Nucleus.


图片



Immunocytochemistry/ Immunofluorescence - Anti-TBLR1/TBL1XR1 antibody [4F3-A8-D9] (ab117761)








All lanes : Anti-TBLR1/TBL1XR1 antibody [4F3-A8-D9] (ab117761) at 1/1000 dilution

Lane 1 : COS-7 (African green monkey kidney fibroblast-like cell line) cell lysate Lane 2 : CEM cell lysate Lane 3 : K562 cell lysate Lane 4 : HeLa cell lysate

Predicted band size: 56 kDa




Western blot - Anti-TBLR1/TBL1XR1 antibody [4F3-A8-D9] (ab117761) All lanes : Anti-TBLR1/TBL1XR1 antibody [4F3-A8-D9] (ab117761) at 1/1000 dilution

Lane 1 : Mouse Brain Lysate Lane 2 : Mouse Heart Lysate Lane 3 : K562 cell Lysate


Predicted band size: 56 kDa

This image was produced with antibody produced in ascites.




ab117761, at 1/500 - 1/1000 dilution, staining TBLR1/TBL1XR1 in paraffin-embedded Human breast cancer by Imunohistochemistry. This image was produced with antibody produced in ascites.

Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-TBLR1/TBL1XR1 antibody [4F3-A8-D9] (ab117761)



Immunocytochemical staining of HeLa cells using anti-TBLR1/TBL1XR1 antibody (ab117761) at a dilution of 1:200. Cells were fixed with 4% paraformaldehyde. This image was produced with antibody produced in ascites.


Immunocytochemistry/ Immunofluorescence - Anti-TBLR1/TBL1XR1 antibody [4F3-A8-D9] (ab117761)



Anti-TBLR1/TBL1XR1 antibody [4F3-A8-D9] (ab117761) at 1/1000 dilution + K562 whole cell lysate at 40 µg

Predicted band size: 56 kDa Observed band size: 60 kDa

This image was produced with antibody produced in ascites.



All lanes : Anti-TBLR1/TBL1XR1 antibody [4F3-A8-D9] (ab117761) at 1/1000 dilution Lane 1 : Mouse heart lysate Lane 2 : Mouse brain lysate Lysates/proteins at 40 µg per lane.

Predicted band size: 56 kDa Observed band size: 60 kDa

A8-D9] (ab117761)

This image was produced with antibody produced in ascites.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you •
- We investigate all quality concerns to ensure our products perform to the highest standards ٠

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

#### Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors