

Anti-TFIIB antibody [EP4588] ab109518

RabMAb

5 图像

概述

产品名称	Anti-TFIIB抗体[EP4588]
描述	兔单克隆抗体[EP4588] to TFIIB
宿主	Rabbit
经测试应用	适用于: WB, IP, IHC-P, Flow Cyt (Intra) 不适用于: ICC/IF
种属反应性	与反应: Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: HepG2, HeLa, Jurkat and K562 cell lysates; IHC-P: Human breast Tissue IP: Jurkat cells Flow Cyt (intra): Jurkat
常规说明	Our RabMAb [®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents . The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing. If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
存储溶液	pH: 7.20 Preservative: 0.05% Sodium azide Constituents: 0.1% BSA, 40% Glycerol (glycerin, glycerine), 9.85% Tris glycine, 50% Tissue culture supernatant
纯度	Tissue culture supernatant
克隆	单克隆

同种型 IgG

The Abpromise guarantee

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/1000 - 1/10000. Predicted molecular weight: 35 kDa.
IP		1/10 - 1/100.
IHC-P		1/100 - 1/250. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol. Antigen retrieval is recommended.
Flow Cyt (Intra)		1/10 - 1/100. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

应用说明 Is unsuitable for ICC/IF.


功能

序列相似性 Belongs to the TFIIIB family.

Contains 1 TFIIIB-type zinc finger.

细胞定位 Nucleus.

图片



All lanes : Anti-TFIIIB antibody [EP4588] (ab109518) at 1/1000 dilution

Lane 1 : C6 (Rat glial tumor glial cell) whole cell lysate

Lane 2 : Raw264.7 (Mouse Abelson murine leukemia virus-induced tumor macrophage) whole cell lysate

Lane 3 : PC-12 (Rat adrenal gland pheochromocytoma) whole cell lysate

Lane 4 : NIH/3T3 (Mouse embryonic fibroblast) whole cell lysate


Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/1000 dilution

Predicted band size: 35 kDa

Observed band size: 35 kDa



ab109518 (purified) at 1/20 dilution (0.16 µg/ml)

immunoprecipitating TFIIIB in Jurkat whole cell lysate .


Lane 1 (input): Jurkat (Human T cell leukemia T lymphocyte) whole cell lysate 10µg

Lane 2 (+): ab109518 & Jurkat whole cell lysate

Lane 3 (-): Rabbit monoclonal IgG ([ab172730](#)) instead of ab109518 in Jurkat whole cell lysate.


For western blotting, ab109518 at 1/500 and VeriBlot for IP Detection Reagent (HRP) ([ab131366](#)) was used for detection at 1/1000 dilution.

Blocking and diluting buffer: 5% NFDM /TBST.



Flow Cytometry (Intracellular) - Anti-TFIIIB antibody
[EP4588] (ab109518)

Intracellular Flow Cytometry analysis of Jurkat (Human T cell leukemia T lymphocyte) cells labelling TFIIIB with ab109518 at 1/20 dilution (4.05 µg/ml) (Red). Cells were fixed with 4% paraformaldehyde. Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) was used as the secondary antibody at 1/2000 dilution. Isotype control - Rabbit monoclonal IgG (**ab172730**) (Black). Unlabeled control - Unlabelled cells (Blue).



Western blot - Anti-TFIIIB antibody [EP4588]
(ab109518)

All lanes : Anti-TFIIIB antibody [EP4588] (ab109518) at 1/1000 dilution

Lane 1 : HepG2 cell lysate


Lane 2 : HeLa cell lysate

Lane 3 : Jurkat cell lysate

Lane 4 : K562 cell lysate

Lysates/proteins at 10 µg per lane.

Predicted band size: 35 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TFIIIB antibody [EP4588] (ab109518)

ab109518, at a 1/100 dilution, staining TFIIIB in paraffin embedded Human breast tissue by Immunohistochemistry.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors