abcam

Product datasheet

Anti-TRAP/CD40L antibody [MR1] ab25206

概述

产品名称 Anti-TRAP/CD40L抗体[MR1]

描述 亚美尼亚仓鼠单克隆抗体[MR1] to TRAP/CD40L

宿主 Armenian hamster

特异性 ab25206 recognises TRAP/CD40L, Mw 39-kDa.

经测试应用 适用于: IHC-Fr, Blocking, IP, Flow Cyt

种属反应性 与反应: Mouse

免疫原 Tissue, cells or virus corresponding to Mouse TRAP/CD40L. Activated mouse Th1 clone D1.6

常规说明 Administration of this antibody to mice blocks the ability to mount primary and secondary immune

responses to TD antigens, yet does not alter the immune response to TI antigens.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

性能

同种型

形式 Liquid

存放说明 Shipped at 4°C. Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.

存储溶液 pH: 8.20

Constituent: 100% Borate buffered saline

纯**度** Affinity purified

Primary antibody说明 Administration of this antibody to mice blocks the ability to mount primary and secondary immune

responses to TD antigens, yet does not alter the immune response to TI antigens.

克隆 单克隆

克隆编号 MR1

轻链类型 kappa

ΙgG

1

The Abpromise guarantee Abpromise Makika Abpromise Maki

Abpromise™承诺保证使用ab25206于以下的经测试应用

"应用说明"部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-Fr		Use at an assay dependent concentration.
Blocking		Use at an assay dependent concentration.
IP		Use at an assay dependent concentration.
Flow Cyt		Use at an assay dependent concentration. <u>ab18479</u> - Armenian Hamster monoclonal lgG, is suitable for use as an isotype control with this antibody.

靶标

功能 Mediates B-cell proliferation in the absence of co-stimulus as well as IgE production in the

presence of IL-4. Involved in immunoglobulin class switching.

Release of soluble CD40L from platelets is partially regulated by GP Ilb/Illa, actin polymerization,

and an matrix metalloproteinases (MMP) inhibitor-sensitive pathway.

组织特异性 Specifically expressed on activated CD4+ T-lymphocytes.

疾病相关 Defects in CD40LG are the cause of X-linked immunodeficiency with hyper-lgM type 1 (HIGM1)

[MIM:308230]; also known as X-linked hyper IgM syndrome (XHIM). HIGM1 is an immunoglobulin isotype switch defect characterized by elevated concentrations of serum IgM and decreased amounts of all other isotypes. Affected males present at an early age (usually within the first year of life) recurrent bacterial and opportunistic infections, including Pneumocystis carinii pneumonia and intractable diarrhea due to cryptosporidium infection. Despite substitution treatment with intravenous immunoglobulin, the overall prognosis is rather poor, with a death rate of about 10%

before adolescence.

序列相似性 Belongs to the tumor necrosis factor family.

翻译后修饰 The soluble form derives from the membrane form by proteolytic processing.

N-linked glycan is a mixture of high mannose and complex type. Glycan structure does not

influence binding affinity to CD40.

Not O-glycosylated.

细**胞定位** Secreted and Cell membrane.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors