

Anti-TSLP antibody ab47943

28 References 2 图像

概述	
产品名称	Anti-TSLP抗体
描述	兔多克隆抗体to TSLP
宿主	Rabbit
经测试应用	适用于: IHC-P, ICC/IF, WB, ELISA, Sandwich ELISA
种属反应性	与反应: Human
免疫原	Recombinant full length TSLP protein (Human),>98% pure
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>
性能	
形式	Lyophilized:Reconstitute with 100ul of sterile water to 1mg/ml.
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
纯度	Immunogen affinity purified
纯化说明	This antibody was purified by affinity chromatography employing immobilized hTSLP matrix.
克隆	多克隆
同种型	IgG
应用	

The Abpromise guarantee **Abpromise™**承诺保证使用ab47943于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-P		Use at an assay dependent concentration.
ICC/IF		Use at an assay dependent concentration.
WB		Use a concentration of 0.1 - 0.2 µg/ml. Detects a band of approximately 28 kDa (predicted molecular weight: 18 kDa). Detection limit for recombinant hTSLP is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.
ELISA		Use at an assay dependent concentration. This allows the detection of 0.2 - 0.4 ng/well of recombinant human TSLP, using 100µl/well antibody solution.
Sandwich ELISA		Use at an assay dependent concentration. To detect hTSLP by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with ab84247 detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hTSLP.


靶标

功能	Cytokine that induces the release of T cell-attracting chemokines from monocytes and, in particular, enhances the maturation of CD11c(+) dendritic cells. Can induce allergic inflammation by directly activating mast cells.
组织特异性	Expressed in a number of tissues including heart, liver and prostate.
细胞定位	Secreted.

图片



ab47943 detecting TSLP in sandwich ELISA



Western blot - Anti-TSLP antibody (ab47943)

Anti-TSLP antibody (ab47943) at 1 µg/ml + Human liver tissue lysate - total protein (**ab29889**) at 10 µg

Secondary

Goat Anti-Rabbit IgG H&L (HRP) preadsorbed (**ab97080**) at 1/5000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 18 kDa

Observed band size: 28 kDa

Additional bands at: 48 kDa, 51 kDa, 53 kDa. We are unsure as to the identity of these extra bands.

Exposure time: 4 minutes

The 28 kDa band observed is comparable to the molecular weight seen with other commercially available antibodies to TSLP.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors