

# Anti-TXNIP antibody [EPR14774] - BSA and Azide free ab215366

敲除验证
重组
RabMAb

[3 References](#)
[4 图像](#)

### 概述

<b>产品名称</b>	Anti-TXNIP抗体[EPR14774] - BSA and Azide free
<b>描述</b>	兔单克隆抗体[EPR14774] to TXNIP - BSA and Azide free
<b>宿主</b>	Rabbit
<b>特异性</b>	<p>Expression levels of the target protein vary with sample type and some optimisation may be required.&amp;nbsp; For western blotting, more concentrated lysates may be required when using tissues samples.</p> <p>Stimulation may be required to allow detection of the target protein due to low levels of endogenous expression in some samples. Please see images below for recommended treatment conditions and positive controls.</p>
<b>经测试应用</b>	<b>适用于:</b> WB, IHC-P
<b>种属反应性</b>	<b>与反应:</b> Mouse, Rat, Human
<b>免疫原</b>	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
<b>阳性对照</b>	Human fetal thymus, BxPC-3, PC-12 and NIH/3T3 lysates; Human kidney and Mouse liver tissues; HeLa and U87-MG cells. WB : Wild-type HAP1 whole cell lysate, MCF7 whole cell lysate and K562 whole cell lysate.
<b>常规说明</b>	<p>ab215366 is the carrier-free version of <a href="#">ab188865</a>.</p> <p>Our <b>carrier-free</b> antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our <b>conjugation kits</b> for antibody conjugates that are ready-to-use in as little as 20 minutes with &lt;1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar<sup>®</sup> Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar<sup>®</sup> is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p>

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information [see here](#).

Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to [RabMAb<sup>®</sup> patents](#).

## 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C. Do Not Freeze.
存储溶液	pH: 7.2 Constituent: PBS
无载体	是
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR14774
同种型	IgG

## 应用

**The Abpromise guarantee**      **Abpromise<sup>™</sup>承诺** 保证使用 ab215366 于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度; 实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/1000. Detects a band of approximately 44 kDa (predicted molecular weight: 44 kDa).
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

## 靶标

功能	May act as an oxidative stress mediator by inhibiting thioredoxin activity or by limiting its bioavailability. Interacts with COPS5 and restores COPS5-induced suppression of CDKN1B stability, blocking the COPS5-mediated translocation of CDKN1B from the nucleus to the cytoplasm. Functions as a transcriptional repressor, possibly by acting as a bridge molecule between transcription factors and corepressor complexes, and over-expression will induce G0/G1 cell cycle arrest. Required for the maturation of natural killer cells.
序列相似性	Belongs to the arrestin family.
翻译后修饰	Ubiquitinated; undergoes polyubiquitination catalyzed by ITCH resulting in proteasomal degradation.

## 图片



**All lanes** : Anti-TXNIP antibody [EPR14774] (**ab188865**) at 1/1000 dilution

**Lane 1** : Wild-type HAP1 whole cell lysate

**Lane 2** : TXNIP knockout HAP1 whole cell lysate

**Lane 3** : MCF7 (human breast adenocarcinoma cell line) whole cell lysate

**Lane 4** : K-562 (human chronic myelogenous leukemia lymphoblasts cell line) whole cell lysate

Lysates/proteins at 20 µg per lane.


**Predicted band size:** 44 kDa

**Observed band size:** 44 kDa

**Lanes 1 - 4:** Merged signal (red and green). Green - **ab188865** observed at 44 kDa. Red - loading control, **ab18058**, observed at 130 kDa.

**ab188865** was shown to recognize TXNIP in wild-type HAP1 cells as signal was lost at the expected MW in TXNIP knockout cells. Additional cross-reactive bands were observed in the wild-type and knockout cells. Wild-type and TXNIP knockout samples were subjected to SDS-PAGE. **ab188865** and **ab18058** (Mouse anti Vinculin loading control) were incubated overnight at 4°C at 1/1000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed **ab216773** and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed **ab216776** secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab188865**).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TXNIP antibody [EPR14774] - BSA and Azide free (ab215366)

Immunohistochemical analysis of paraffin-embedded Human kidney tissue labeling TXNIP with [ab188865](#) at 1/250 dilution followed by prediluted HRP Polymer for Rabbit IgG. Counter stained with Hematoxylin.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab188865](#)).

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TXNIP antibody [EPR14774] - BSA and Azide free (ab215366)

Immunohistochemical analysis of paraffin-embedded Mouse liver tissue labeling TXNIP with [ab188865](#) at 1/250 dilution followed by prediluted HRP Polymer for Rabbit IgG. Counter stained with Hematoxylin.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab188865](#)).

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

### Why choose a recombinant antibody?



**Research with confidence**  
Consistent and reproducible results



**Long-term and scalable supply**  
Recombinant technology



**Success from the first experiment**  
Confirmed specificity



**Ethical standards compliant**  
Animal-free production

Anti-TXNIP antibody [EPR14774] - BSA and Azide free (ab215366)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors