


Anti-URP2/Kindlin-3 antibody ab68040

★★★★★ 6 Abreviews 10 References 2 图像

概述	
产品名称	Anti-URP2/Kindlin-3抗体
描述	兔多克隆抗体to URP2/Kindlin-3
宿主	Rabbit
特异性	From Jan 2024, QC testing of replenishment batches of this polyclonal changed. All tested and expected application and reactive species combinations are still covered by our Abcam product promise. However, we no longer test all applications. For more information on a specific batch, please contact our Scientific Support who will be happy to help.
经测试应用	适用于: IHC-P, WB
种属反应性	与反应: Mouse, Rat, Human 预测可用于: Cow 
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>
性能	
形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	pH: 7.40 Preservative: 0.02% Sodium azide Constituent: PBS
Batches of this product that have a concentration < 1mg/ml may have BSA added as a stabilising agent. If you would like information about the formulation of a specific lot, please contact our scientific support team who will be happy to help.	

纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab68040于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-P		Use a concentration of 5 µg/ml.
WB	★★★★★ (2)	Use a concentration of 1 µg/ml. Detects a band of approximately 76 kDa (predicted molecular weight: 76 kDa).

靶标

功能	Plays a central role in cell adhesion in hematopoietic cells. Acts by activating the integrin beta-1-3 (ITGB1, ITGB2 and ITGB3). Required for integrin-mediated platelet adhesion and leukocyte adhesion to endothelial cells. Required for activation of integrin beta-2 (ITGB2) in polymorphonuclear granulocytes (PMNs). Isoform 2 may act as a repressor of NF-kappa-B and apoptosis.
组织特异性	Highly expressed in lymph node. Expressed in thymus, spleen and leukocytes. Weakly expressed in placenta, small intestine, stomach, testis and lung. Overexpressed in B-cell malignancies.
疾病相关	Defects in FERMT3 are the cause of leukocyte adhesion deficiency type 3 (LAD3) [MIM:612840]; also called leukocyte adhesion deficiency 1 variant (LAD1v). LAD3 is a rare syndrome characterized by infections without pus formation in the presence of a leukocytosis combined with a Glanzmann-type bleeding disorder, resulting from a hematopoietic defect in integrin activation. Symptoms arise from an inability to activate the integrins expressed on hematopoietic cells, including platelets and leukocytes.
序列相似性	Belongs to the kindlin family. Contains 1 FERM domain. Contains 1 PH domain.
结构域	The FERM domain is not correctly detected by PROSITE or Pfam techniques because it contains the insertion of a PH domain.
细胞定位	Cell projection > podosome. Present in the F-actin surrounding ring structure of podosomes, which are specialized adhesion structures of hematopoietic cells.

图片



Western blot - Anti-URP2/Kindlin-3 antibody
(ab68040)

All lanes : Anti-URP2/Kindlin-3 antibody (ab68040) at 1 µg/ml

Lane 1 : Kidney (Rat) Tissue Lysate

Lane 2 : Spleen (Mouse) Tissue Lysate

Lane 3 : Thymus (Rat) Tissue Lysate

Lysates/proteins at 10 µg/ml per lane.

Secondary


All lanes : Goat polyclonal to Rabbit IgG - H&L - Pre-Adsorbed (HRP) at 1/3000 dilution

Performed under reducing conditions.

Predicted band size: 76 kDa

Observed band size: 76 kDa

Exposure time: 2 minutes



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-URP2/Kindlin-3 antibody
(ab68040)

IHC image of URP2/Kindlin-3 staining in Human normal lung formalin fixed paraffin embedded tissue section, performed on a Leica Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab68040, 5µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors