abcam

Product datasheet

Anti-VEGF Receptor 2 antibody ab39638

★★★★★ <u>3 Abreviews</u> <u>21 References</u> 6 图像

概述		
产品名称	Anti-VEGF Receptor 2 抗体	
描述	兔多克隆抗体to VEGF Receptor 2	
宿主	Rabbit	
经 测 试应 用	适用于: ICC/IF, WB, IHC-P	
种属反 应性	与反应: Rat, Human	
	预测可用于: Mouse 🔺	
免疫原	The antiserum was produced against synthesized non-phosphopeptide derived from human VEGF Receptor 2 around the phosphorylation site of tyrosine 951.	
阳性 对照	WB: SK-OV3 cell lysate. IHC-P: Human brain tissue and stomach cancer tissue; Rat spinal cord tissue. ICC/IF: HepG2 cells.	
常 规说 明	The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.	
	If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As	
性能		
形式	Liquid	
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.	
存储溶液	pH: 7.40 Preservative: 0.02% Sodium azide	

克隆	多克隆
纯 化说明	This antibody was affinity purified from rabbit antiserum by affinity chromatography using epitope specific immunogen.
纯 度	Immunogen affinity purified
	Without Mg2+ and Ca2+
	Constituents: PBS, 50% Glycerol, 0.87% Sodium chloride
	Preservative: 0.02% Sodium azide

应用


The Abpromise guarantee Abpromise™承诺保证使用ab39638于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC/IF	★★★★ ★ (<u>1)</u>	Use a concentration of 1 µg/ml.
WB	★★★★★ <u>(1)</u>	1/500 - 1/1000. Predicted molecular weight: 151 kDa.
IHC-P	★ ★ ★ ★ ★ (1)	1/50 - 1/100.


靶 标	
功能	Receptor for VEGF or VEGFC. Has a tyrosine-protein kinase activity. The VEGF-kinase ligand/receptor signaling system plays a key role in vascular development and regulation of vascular permeability. In case of HIV-1 infection, the interaction with extracellular viral Tat protein seems to enhance angiogenesis in Kaposi's sarcoma lesions.
疾病相关	Defects in KDR are associated with susceptibility to hemangioma capillary infantile (HCI) [MIM:602089]. HCI are benign, highly proliferative lesions involving aberrant localized growth of capillary endothelium. They are the most common tumor of infancy, occurring in up to 10% of all births. Hemangiomas tend to appear shortly after birth and show rapid neonatal growth for up to 12 months characterized by endothelial hypercellularity and increased numbers of mast cells. This phase is followed by slow involution at a rate of about 10% per year and replacement by fibrofatty stroma.
序列相似性	Belongs to the protein kinase superfamily. Tyr protein kinase family. CSF-1/PDGF receptor subfamily. Contains 7 lg-like C2-type (immunoglobulin-like) domains. Contains 1 protein kinase domain.
翻 译后修饰	Phosphorylated. Dephosphorylated by PTPRB. Dephosphorylated by PTPRJ at Tyr-951, Tyr-996, Tyr-1054, Tyr-1059, Tyr-1175 and Tyr-1214.
细 胞定位	Membrane.

图片




Western blot analysis of extracts from SK-OV3 cells using VEGF Receptor 2 antibody ab39638 (Lane 1 and 2) and VEGF Receptor 2 (phospho-Tyr951) antibody (Lane 3 and 4).

Secondary antibody - anti-rabbit HRP (ab6721)




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-VEGF Receptor 2 antibody (ab39638)

Immunohistochemical analysis of human brain tissue stained with ab39638. The image on the right is blocked with the synthesized peptide.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-VEGF Receptor 2 antibody (ab39638)




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-VEGF Receptor 2 antibody (ab39638)

Immunocytochemistry/ Immunofluorescence - Anti-VEGF Receptor 2 antibody (ab39638) Immunohistochemical analysis of human stomach cancer tissue stained with ab39638 at 1/200 dilution at 4°C overnight. The negative control was secondary antibody only.

Immunohistochemical analysis of rat spinal cord tissue stained with ab39638 at 1/200 dilution at 4°C overnight. The negative control was secondary antibody only.

ICC/IF image of ab39638 stained HepG2 cells. The cells were 4% PFA fixed (10 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab39638, 1 μ g/ml) overnight at +4°C. The secondary antibody

(green) was Alexa Fluor® 488 goat anti-rabbit lgG (H+L) used at a 1/1000 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.



Anti-VEGF Receptor 2 antibody (ab39638) (Anti-VEGF Receptor 2 antibody) + HeLa cell lysate

Predicted band size: 151 kDa

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors