abcam

Product datasheet

Anti-ZNF787 antibody ab117837

2 图像

概述

免疫原

产**品名称** Anti-ZNF787抗体

描述 兔多克隆抗体to ZNF787

宿主 Rabbit

经测试应用 适用于: WB, IP

种属反应性 与反应: Mouse, Human

预测可用于: Rat, Guinea pig, Cow, Dog, Pig, Rhesus monkey, Gorilla, Elephant

Synthetic peptide, corresponding to a region within amino acids 1-50 of Human ZNF787

(NP_001002836.2).

阳性对照 HeLa, 293T, Jurkat and Mouse NIH3T3 whole cell lysates.

常规说明

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

性能

形式 Liquid

存放说明 Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

存储溶液 pH: 6.8

Preservative: 0.09% Sodium azide

Constituents: 99% Tris buffered saline, 0.1% BSA

纯**度** Immunogen affinity purified

纯**化说明** ab117837 was affinity purified using an epitope specific to ZNF787 immobilized on solid support.

克隆 多克隆

同种型 IgG

The Abpromise guarantee

Abpromise™承诺保证使用ab117837于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/2000 - 1/10000. Predicted molecular weight: 40 kDa.
IP		Use at 2-10 μg/mg of lysate.

靶标

功能 May be involved in transcriptional regulation.


序列相似性 Belongs to the krueppel C2H2-type zinc-finger protein family.

Contains 7 C2H2-type zinc fingers.

翻译后修饰 Phosphorylated upon DNA damage, probably by ATM or ATR.

细胞定位 Nucleus.

图片



Western blot - Anti-ZNF787 antibody (ab117837)

All lanes: Anti-ZNF787 antibody (ab117837) at 0.04 µg/ml


Lane 1 : HeLa whole cell lysate at 50 μg Lane 2 : HeLa whole cell lysate at 15 μg Lane 3 : 293T whole cell lysate at 50 μg Lane 4 : Jurkat whole cell lysate at 50 μg

Lane 5: NIH3T3 whole cell lysate at 50 µg

Developed using the ECL technique.

Predicted band size: 40 kDa

Exposure time: 3 minutes



Detection of Human ZNF787 by Immunoprecipitation in HeLa whole cell lysate (1 mg for IP, 20% of IP loaded) using ab117837 at 6 μ g/mg of lysate. Subsequent WB detection was done using ab117837 at 1 μ g/ml. ECL detection. Exposure time: 3 seconds.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors