




Anti-ZRANB1 antibody ab82132

1 References **1 图像**

概述

产品名称	Anti-ZRANB1抗体
描述	山羊多克隆抗体to ZRANB1
宿主	Goat
经测试应用	适用于: WB
种属反应性	与反应: Human 预测可用于: Mouse, Cow 
免疫原	Synthetic peptide: KWLDRYRQIRPCTS , corresponding to C terminal amino acids 682-695 of Human ZRANB1  Run BLAST with  Run BLAST with
阳性对照	Human Cerebellum Lysate
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	pH: 7.30 Preservative: 0.02% Sodium azide Constituents: 0.5% BSA, Tris buffered saline
纯度	Immunogen affinity purified
纯化说明	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
克隆	多克隆

同种型

IgG

应用

The Abpromise guarantee

Abpromise™承诺保证使用ab82132于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		Use a concentration of 1 - 3 µg/ml. Predicted molecular weight: 81 kDa. 1 hour primary incubation is recommended for this product.

靶标

功能	Positive regulator of the Wnt signaling pathway that specifically cleaves 'Lys-63'-linked ubiquitin chains. Acts by deubiquitinating APC protein, a negative regulator of Wnt-mediated transcription. May also modulate TNF-alpha signaling.
组织特异性	Widely expressed.
序列相似性	Belongs to the peptidase C64 family. Contains 1 OTU domain. Contains 3 RanBP2-type zinc fingers.
结构域	The OTU domain mediates the deubiquitinating activity. The RanBP2-type zinc fingers mediate the specific interaction with 'Lys-63'-linked ubiquitin.
细胞定位	Cytoplasm. Nucleus.

图片



Anti-ZRANB1 antibody (ab82132) at 0.1 µg/ml + Human Cerebellum Lysate (35ug in RIPA buffer)

Predicted band size: 81 kDa

Primary incubation was 1 hour. Detected by chemiluminescence.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors