

ALK Mutant (T1151_L1152insT), Active ab186465

5 图像

描述

产品名称	ALK Mutant (T1151_L1152insT), Active
生物活性	Specific activity of ab186465 was determined to be 13 nmol/min/mg as per activity assay protocol.
纯度	> 80 % Densitometry. Affinity purified.
表达系统	Insect cells
Accession	<u>Q9UM73</u>
蛋白长度	Protein fragment
无动物成分	No
性质	Recombinant
种属	Human
序列	<pre> RRKHQELQAMQMELQSPEYKLSKLRSTIMTDYNPNYCFAGK TSSISDLK EVPRKNITLIRGLGHGAFGEVYEGQVSGMPNDPSPLQVAVKL PEVCSEQD ELDFLMEALIISKFNHQNIVRCIGVSLQSLPRFILLELMAGG DLKSFLRE TRPRPSQPSSLAMLDLLHWARDIACGCQYLEENHFIIHRDIAA RNCLLTCP GPGRVAKIGDFGMARDIYRASYYRKGGCAMLVWKMPPEAFM EGIFTSKT DTWSFGVLLWEIFSLGYMPYPSKSNQEVLEFVTSGGRMDPPK NCPGPVYR IMTQCWQHQPEDRPNFAIILERIEYCTQDPDVINTALPIEYG PLVEEEEK VPVRPKDPEGVPPLLVSQQAQKREERSPAAPPPLPTTSSGKA AKKPTAAE ISVRVPRGPAVEGGHVNMAFSQSNPPSELHKVHGSRNKPTSL WNPTYGSW FTEKPTKKNPIAKKEPHDRGNLGLGEGSCTVPPNVATGRLPG ASLLEPS SLTANMKEVPLFRLRHFP CGNVNYGYQQGLPLEAATAPGAG HYEDTILK SKNSMNQPGP </pre>

预测分子量	90 kDa including tags
氨基酸	1060 to 1620
标签	GST tag N-Terminus
额外的序列信息	Cytoplasmic domain ALK Mutant (T1151_L1152insT)
描述	重组人ALK蛋白

技术指标

Our **Abpromise guarantee** covers the use of **ab186465** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	SDS-PAGE Functional Studies
形式	Liquid

制备和贮存

稳定性和存储	Shipped on Dry Ice. Upon delivery aliquot. Store at -80°C. Avoid freeze / thaw cycle. pH: 7.5 Preservative: 1.02% Imidazole Constituents: 0.82% Sodium phosphate, 1.74% Sodium chloride, 0.002% PMSF, 0.004% DTT, 25% Glycerol (glycerin, glycerine) This product is an active protein and may elicit a biological response in vivo, handle with caution.
--------	---

常规信息

功能	Neuronal receptor tyrosine kinase that is essentially and transiently expressed in specific regions of the central and peripheral nervous systems and plays an important role in the genesis and differentiation of the nervous system. Transduces signals from ligands at the cell surface, through specific activation of the mitogen-activated protein kinase (MAPK) pathway. Phosphorylates almost exclusively at the first tyrosine of the Y-x-x-x-Y-Y motif. Following activation by ligand, ALK induces tyrosine phosphorylation of CBL, FRS2, IRS1 and SHC1, as well as of the MAP kinases MAPK1/ERK2 and MAPK3/ERK1. Acts as a receptor for ligands pleiotrophin (PTN), a secreted growth factor, and midkine (MDK), a PTN-related factor, thus participating in PTN and MDK signal transduction. PTN-binding induces MAPK pathway activation, which is important for the anti-apoptotic signaling of PTN and regulation of cell proliferation. MDK-binding induces phosphorylation of the ALK target insulin receptor substrate (IRS1), activates mitogen-activated protein kinases (MAPKs) and PI3-kinase, resulting also in cell proliferation induction. Drives NF-kappa-B activation, probably through IRS1 and the activation of the AKT serine/threonine kinase. Recruitment of IRS1 to activated ALK and the activation of NF-kappa-B are essential for the autocrine growth and survival signaling of MDK.
组织特异性	Expressed in brain and CNS. Also expressed in the small intestine and testis, but not in normal lymphoid cells.
疾病相关	A chromosomal aberration involving ALK is found in a form of non-Hodgkin lymphoma. Translocation t(2;5)(p23;q35) with NPM1. The resulting chimeric NPM1-ALK protein homodimerize and the kinase becomes constitutively activated. The constitutively active fusion proteins are responsible for 5-10% of non-Hodgkin lymphomas.

A chromosomal aberration involving ALK is associated with inflammatory myofibroblastic tumors (IMTs). Translocation t(2;11)(p23;p15) with CARS; translocation t(2;4)(p23;q21) with SEC31A. A chromosomal aberration involving ALK is associated with anaplastic large-cell lymphoma (ALCL). Translocation t(2;17)(p23;q25) with ALO17.

Neuroblastoma 3

The ALK signaling pathway plays an important role in glioblastoma, the most common malignant brain tumor of adults and one of the most lethal cancers. It regulates both glioblastoma migration and growth.

A chromosomal aberration involving ALK is found in one subject with colorectal cancer. Translocation t(2;2)(p23.1;p23.3). A 5 million base pair tandem duplication generates an in-frame WDCP-ALK gene fusion.

序列相似性

Belongs to the protein kinase superfamily. Tyr protein kinase family. Insulin receptor subfamily. Contains 1 LDL-receptor class A domain. Contains 2 MAM domains. Contains 1 protein kinase domain.


翻译后修饰

Phosphorylated at tyrosine residues by autocatalysis, which activates kinase activity. In cells not stimulated by a ligand, receptor protein tyrosine phosphatase beta and zeta complex (PTPRB/PTPRZ1) dephosphorylates ALK at the sites in ALK that are undergoing autophosphorylation through autoactivation. Phosphorylation at Tyr-1507 is critical for SHC1 association. N-glycosylated.

细胞定位


Cell membrane. Membrane attachment was crucial for promotion of neuron-like differentiation and cell proliferation arrest through specific activation of the MAP kinase pathway.

图片




The specific activity of ALK (ab186465) was determined to be 49 nmol/min/mg as per activity assay protocol and was equivalent to 200 nmol/min/mg as per radiometric assay

Functional Studies - ALK Mutant
(T1151_L1152insT), Active (ab186465)




SDS PAGE analysis of ab186465

SDS-PAGE - ALK Mutant (T1151_L1152insT),
Active (ab186465)




SDS PAGE analysis of ab186465

SDS-PAGE - ALK Mutant (T1151_L1152insT),
Active (ab186465)



Specific activity of ab186465 was determined to be 13 nmol/min/mg as per activity assay protocol.

Functional Studies - ALK Mutant
(T1151_L1152insT), Active (ab186465)



Purity of ab186465 was determined to be >80% by densitometry.
Approx. MW 90kDa.

SDS-PAGE - ALK Mutant (T1151_L1152insT),
Active (ab186465)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors