abcam

Product datasheet

Lipocalin-2 / NGAL peptide ab187960

1 图像

描述

产**品名称** Lipocalin-2 / NGAL多肽

 Accession
 P80188

 无动物成分
 No

性质 Synthetic

技术指标

应用

Our Abpromise quarantee covers the use of ab187960 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

形式 Liquid

补充说明 - First try to dissolve a small amount of peptide in either water or buffer. The more charged

residues on a peptide, the more soluble it is in aqueous solutions.

- If the peptide doesn't dissolve try an organic solvent e.g. DMSO, then dilute using water or

Blocking - Blocking peptide for Anti-Lipocalin-2 / NGAL antibody [EPR5084] (ab125075)

buffer.

- Consider that any solvent used must be compatible with your assay. If a peptide does not

dissolve and you need to recover it, lyophilise to remove the solvent.

- Gentle warming and sonication can effectively aid peptide solubilisation. If the solution is

cloudy or has gelled the peptide may be in suspension rather than solubilised.

- Peptides containing cysteine are easily oxidised, so should be prepared in solution just prior

to use.

制备和贮存

稳定性和存储 Shipped at 4°C. Store at -20°C.

常规信息

功能 Iron-trafficking protein involved in multiple processes such as apoptosis, innate immunity and

 $renal\ development.\ Binds\ iron\ through\ association\ with\ 2,5-dihydroxybenzoic\ acid\ (2,5-DHBA),\ a$ $side rophore\ that\ shares\ structural\ similarities\ with\ bacterial\ enterobactin,\ and\ delivers\ or\ removes$

iron from the cell, depending on the context. Iron-bound form (holo-24p3) is internalized following

binding to the SLC22A17 (24p3R) receptor, leading to release of iron and subsequent increase of intracellular iron concentration. In contrast, association of the iron-free form (apo-24p3) with the SLC22A17 (24p3R) receptor is followed by association with an intracellular siderophore, iron chelation and iron transfer to the extracellular medium, thereby reducing intracellular iron concentration. Involved in apoptosis due to interleukin-3 (IL3) deprivation: iron-loaded form increases intracellular iron concentration without promoting apoptosis, while iron-free form decreases intracellular iron levels, inducing expression of the proapoptotic protein BCL2L11/BIM, resulting in apoptosis. Involved in innate immunity, possibly by sequestrating iron, leading to limit bacterial growth.

组织特异性

Expressed in bone marrow and in tissues that are prone to exposure to microorganism. High expression is found in bone marrow as well as in uterus, prostate, salivary gland, stomach, appendix, colon, trachea and lung. Not found in the small intestine or peripheral blood leukocytes.


序列相似性

Belongs to the calycin superfamily. Lipocalin family.

细胞定位

Secreted. Upon binding to the SLC22A17 (24p3R) receptor, it is internalized.

图片



To learn more about our protein and peptide range click **here**.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors