

Recombinant Human 4E-BP2 protein ab104667

1 References 1 图像

描述	
产品名称	重组人4E-BP2蛋白
纯度	> 85 % SDS-PAGE. purified by using anion-exchange chromatography (DEAE sepharose resin) and gel-filtration chromatography (Sephacryl S-200) with 20mM Tris pH 7.5, 2mM EDTA.
表达系统	Escherichia coli
Accession	<u>Q13542</u>
蛋白长度	Full length protein
无动物成分	No
性质	Recombinant
种属	Human
序列	MGSSHHHHHHSSGLVPRGSHMSSSAGSGHQPSQSRAIPTR TVAISDAAQL PHDYCTTPGGTLFSTTPGGTRIIYDRKFLLDRRNSPMAQTPP CHLPNIPG VTSPGTLIEDSKVEVNNLNNLNNHDKHAVGDDAQFEMDI
预测分子量	15 kDa including tags
氨基酸	1 to 120
标签	His tag N-Terminus
技术指标	
Our <b>Abpromise guarantee</b> covers the use of <b>ab104667</b> in the following tested applications.	
The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.	
应用	SDS-PAGE Mass Spectrometry
质谱法	MALDI-TOF
形式	Liquid
制备和贮存	
稳定性和存储	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw

cycles.

pH: 8.00

Constituents: 0.0154% DTT, 0.316% Tris HCl, 10% Glycerol (glycerin, glycerine), 0.58% Sodium chloride

## 常规信息

---

### 功能

Repressor of translation initiation involved in synaptic plasticity, learning and memory formation (By similarity). Regulates EIF4E activity by preventing its assembly into the eIF4F complex: hypophosphorylated form of EIF4EBP2 competes with EIF4G1/EIF4G3 and strongly binds to EIF4E, leading to repress translation. In contrast, hyperphosphorylated form dissociates from EIF4E, allowing interaction between EIF4G1/EIF4G3 and EIF4E, leading to initiation of translation (PubMed:25533957). EIF4EBP2 is enriched in brain and acts as a regulator of synapse activity and neuronal stem cell renewal via its ability to repress translation initiation (By similarity). Mediates the regulation of protein translation by hormones, growth factors and other stimuli that signal through the MAP kinase and mTORC1 pathways.

### 序列相似性

Belongs to the eIF4E-binding protein family.

### 结构域


The TOS motif mediates interaction with RPTOR, leading to promote phosphorylation by mTORC1 complex. Intrinsically disordered protein that undergoes folding upon phosphorylation (PubMed:25533957). Hypophosphorylated form interacts strongly with EIF4E using (1) the YXXXXLPhi motif, that undergoes a disorder-to-helix transition upon binding and (2) the secondary EIF4E binding sites (residues 78-82) (PubMed:24207126, PubMed:25533957). Phosphorylation at Thr-37 and Thr-46 induces folding of region encompassing residues from Pro-18 to Arg-62 of into a four-stranded beta-domain that sequesters the helical YXXXXLPhi motif into a buried beta-strand, blocking accessibility to EIF4E. Protein phosphorylated at Thr-37 and Thr-46 is however unstable and subsequent phosphorylation at Ser-65, Thr-70 and Ser-83 is required to stabilize the fold, decreasing affinity for EIF4E by a factor of 4000 (PubMed:24207126, PubMed:25533957).

### 翻译后修饰

Phosphorylation at Thr-37, Thr-46, Ser-65, Thr-70 and Ser-83 is mediated by MTOR and corresponds to the hyperphosphorylated form: it abolishes binding to EIF4E by inducing folding of intrinsically disordered regions (PubMed:24207126, PubMed:25533957). First phosphorylated at Thr-37 and Thr-46 by MTOR, inducing folding of region encompassing residues from Pro-18 to Arg-62 of into a four-stranded beta-domain that sequesters the helical YXXXXLPhi motif into a partly buried beta-strand, blocking accessibility to EIF4E. Protein phosphorylated at Thr-37 and Thr-46 is however unstable and subsequent phosphorylation at Ser-65, Thr-70 and Ser-83 is required to stabilize the fold, decreasing affinity for EIF4E by a factor of 4000 (PubMed:24207126, PubMed:25533957). Phosphorylated in response to insulin, EGF and PDGF. Deamidated at Asn-99 and Asn-102 to aspartate (Asp) in brain. Deamidation promotes interaction with RPTOR, subsequent phosphorylation by mTORC1 and increased translation, leading to impair kinetics of excitatory synaptic transmission. Deamidation takes place during postnatal development, when the PI3K-Akt-mTOR signaling is reduced, suggesting it acts as a compensatory mechanism to promote translation despite attenuated PI3K-Akt-mTOR signaling in neuron development. Deamidation converts Asn residues into a mixture of Asp and isoaspartate; interactions with PCMT1 is required to prevent isoaspartate accumulation and convert isoaspartate to Asp.

## 图片

---



SDS-PAGE of ab104667 (3 $\mu$ g) under reducing condition and visualized by coomassie blue stain.

SDS-PAGE - Recombinant Human 4E-BP2 protein  
(ab104667)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

#### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors