abcam

Product datasheet

Recombinant human ADA protein ab109839

3 References 1 图像

描述

产品名称
重组人ADA蛋白

生物活性 Specific activity is >40 units/mg, and is defined as the amount of enzyme that convert 1.0 umol of

adenosine to inosine per minute at pH 7.5 at 25°C.

纯**度** > 90 % SDS-PAGE.

No detergents such as urea, Triton or Tween were used to purify the proteins. Purified by using anion exchange chromatography (DEAE sepharose resin) and gel filtration chromatography

(Sephacryl S 200) with 20mM Tris pH 7.5, 2mM EDTA.

内毒素水平 < 1.000 Eu/mg

表达系统 Escherichia coli

Accession P00813

蛋白长度 Full length protein

无动物成分 No

性质 Recombinant

种属 Human

序列 MGSSHHHHHH SSGLVPRGSH MAQTPAFDKP

NTAEGLLNVI GMDKPLTLPD FLAKFDYYMP AIAGCREAIK RIAYEFVEMK AKEGVVYVEV RYSPHLLANS KVEPIPWNQA EGDLTPDEVV ALVGQGLQEG ERDFGVKARS ILCCMRHQPN

KVELHVHLDG SIKPETILYY GRRRGIALPA

WSPKVVELCK KYQQQTVVAI DLAGDETIPG SSLLPGHVQA YQEAVKSGIH RTVHAGEVGS AEVVKEAVDI LKTERLGHGY HTLEDQALYN

RLRQENMHFE ICPWSSYLTG AWKPDTEHAV IRLKNDQANY SLNTDDPLIF KSTLDTDYQM TKRDMGFTEE EFKRLNINAA KSSFLPEDEK

RELLDLLYKA YGMPPSASAG QNL

预**测分子量** 43 kDa **氨基酸** 1 to 363

标签 His tag N-Terminus

技术指标

1

Our Abpromise quarantee covers the use of ab109839 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用 Functional Studies

SDS-PAGE

Mass Spectrometry

质**谱法** MALDI-TOF

形式 Liquid

制备和贮存

稳定性和存储 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -

80°C. Avoid freeze / thaw cycle.

pH: 8.00

Constituents: 0.02% DTT, 0.32% Tris HCI, 20% Glycerol (glycerin, glycerine)

This product is an active protein and may elicit a biological response in vivo, handle with caution.

常规信息

功能 Catalyzes the hydrolytic deamination of adenosine and 2-deoxyadenosine. Plays an important

role in purine metabolism and in adenosine homeostasis. Modulates signaling by extracellular adenosine, and so contributes indirectly to cellular signaling events. Acts as a positive regulator of T-cell coactivation, by binding DPP4. Its interaction with DPP4 regulates lymphocyte-epithelial cell

adhesion.

组织**特异性** Found in all tissues, occurs in large amounts in T-lymphocytes and, at the time of weaning, in

gastrointestinal tissues.

疾病相关 Defects in ADA are the cause of severe combined immunodeficiency autosomal recessive T-cell-

[MIM:102700]. SCID refers to a genetically and clinically heterogeneous group of rare congenital disorders characterized by impairment of both humoral and cell-mediated immunity, leukopenia, and low or absent antibody levels. Patients with SCID present in infancy with recurrent, persistent infections by opportunistic organisms. The common characteristic of all types of SCID is absence of T-cell-mediated cellular immunity due to a defect in T-cell development. ADA-SCID is an autosomal recessive form accounting for about 50% of non-X-linked SCIDs. ADA deficiency has

negative/B-cell-negative/NK-cell-negative due to adenosine deaminase deficiency (ADASCID)

been diagnosed in chronically ill teenagers and adults (late or adult onset). Population and newborn screening programs have also identified several healthy individuals with normal immunity


who have partial ADA deficiency.

序列相似性 Belongs to the adenosine and AMP deaminases family.

细胞定位 Cell membrane. Cell junction. Cytoplasmic vesicle lumen. Cytoplasm. Colocalized with DPP4 at

the cell junction in lymphocyte-epithelial cell adhesion.

图片



SDS-PAGE analysis of ab109839 at 3µg. Gel concentration: 15%.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors