

Recombinant human EGFR protein (Fc Chimera) ab84002

2 图像

描述	
产品名称	重组人EGFR蛋白(Fc Chimera)
生物活性	The ED ₅₀ of ab84002 is typically 60-100 ng/ml as measured by its ability to neutralize EGF mediated proliferation of murine NIH3T3 fibroblasts.
纯度	> 95 % SDS-PAGE.
表达系统	HEK 293 cells
Accession	<u>P00533</u>
蛋白长度	Protein fragment
无动物成分	No
性质	Recombinant
种属	Human
序列	<p>Theoretical sequence:</p> <p>LEEKKVCQGTSNKLTLQGTFFEDHFLSLQRMFNCEVVLG NLEITYVQR NYDLSFLKTIQEVAGYVLIALNTVERIPLNLQIIRGNM YYENSYALA VLSNYDANKTGLKELPMRNLQEILHGAVRFSNNPALCNV ESIQWRDIV SSDFLSNMSMDFQNLHGSCQKCDPSCPNGSCWGAGEENC QKLTKIICA QQCSGRCRGKSPSDCCHNQCAAGCTGPRESCLVCRKFR DEATCKDTC PPLMLYNPTTYQMDVNPEGKYSFGATCVKKCPRNYVVT HGSCVRACG ADSYEMEEDGVRKCKCEGPCRKVCNGIGIGEFKDSL NATNIKHF NCTISGDLHILPVAFRGDSFHTPLDLPQELDILKTVK EITGFLLIQ AWPENRTDLHAFENLEIIRGRKQHGQFSLAVVSLNITS LGLRSLKEI SDGDVVIISGNKNLCYANTINWKKLFGTSGQKTKIISNRG ENSCKATGQ VCHALCSPEGCWGPEPRDCVSRSSNTKVDKKVEPKSCDK THTCPPCPA</p>

PELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHED
 PEVKFNWYV
 DGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKE
 YKCKVSNKA
 LPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLT
 CLVKGFYPS
 DIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDK
 SRWQQGNVF SCSVMHEALHNHYTQKSLSLSPGK

氨基酸 1 to 525

额外的序列信息 Fused with the Fc region of Human IgG1 at the C-terminus.

技术指标

Our **Abpromise guarantee** covers the use of **ab84002** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用 SDS-PAGE
 Functional Studies

形式 Lyophilized

制备和贮存

稳定性和存储 Shipped at 4°C. After reconstitution store at -20°C. Avoid freeze / thaw cycles.
 Constituents: 1% Human serum albumin, 10% Trehalose
 This product is an active protein and may elicit a biological response in vivo, handle with caution.

复溶 It is recommended that 0.5 ml of sterile phosphate-buffered saline be added to the vial. Following reconstitution short-term storage at 4°C is recommended, with longer-term storage in aliquots at -18 to -20°C. Repeated freeze thawing is not recommended.

常规信息

功能 Receptor tyrosine kinase binding ligands of the EGF family and activating several signaling cascades to convert extracellular cues into appropriate cellular responses. Known ligands include EGF, TGFA/TGF-alpha, amphiregulin, epigen/EPGN, BTC/betacellulin, epiregulin/EREG and HBEGF/heparin-binding EGF. Ligand binding triggers receptor homo- and/or heterodimerization and autophosphorylation on key cytoplasmic residues. The phosphorylated receptor recruits adapter proteins like GRB2 which in turn activates complex downstream signaling cascades. Activates at least 4 major downstream signaling cascades including the RAS-RAF-MEK-ERK, PI3 kinase-AKT, PLCgamma-PKC and STATs modules. May also activate the NF-kappa-B signaling cascade. Also directly phosphorylates other proteins like RGS16, activating its GTPase activity and probably coupling the EGF receptor signaling to the G protein-coupled receptor signaling. Also phosphorylates MUC1 and increases its interaction with SRC and CTNNB1/beta-catenin.
 Isoform 2 may act as an antagonist of EGF action.

组织特异性 Ubiquitously expressed. Isoform 2 is also expressed in ovarian cancers.

疾病相关 Lung cancer
 Inflammatory skin and bowel disease, neonatal, 2

序列相似性

Belongs to the protein kinase superfamily. Tyr protein kinase family. EGF receptor subfamily. Contains 1 protein kinase domain.

翻译后修饰

Phosphorylation at Ser-695 is partial and occurs only if Thr-693 is phosphorylated. Phosphorylation at Thr-678 and Thr-693 by PRKD1 inhibits EGF-induced MAPK8/JNK1 activation. Dephosphorylation by PTPRJ prevents endocytosis and stabilizes the receptor at the plasma membrane. Autophosphorylation at Tyr-1197 is stimulated by methylation at Arg-1199 and enhances interaction with PTPN6. Autophosphorylation at Tyr-1092 and/or Tyr-1110 recruits STAT3. Dephosphorylated by PTPN1 and PTPN2. Monoubiquitinated and polyubiquitinated upon EGF stimulation; which does not affect tyrosine kinase activity or signaling capacity but may play a role in lysosomal targeting. Polyubiquitin linkage is mainly through 'Lys-63', but linkage through 'Lys-48', 'Lys-11' and 'Lys-29' also occurs. Deubiquitination by OTUD7B prevents degradation. Ubiquitinated by RNF115 and RNF126. Methylated. Methylation at Arg-1199 by PRMT5 stimulates phosphorylation at Tyr-1197.


细胞定位

Secreted and Cell membrane. Endoplasmic reticulum membrane. Golgi apparatus membrane. Nucleus membrane. Endosome. Endosome membrane. Nucleus. In response to EGF, translocated from the cell membrane to the nucleus via Golgi and ER. Endocytosed upon activation by ligand. Colocalized with GPER1 in the nucleus of estrogen agonist-induced cancer-associated fibroblasts (CAF).

图片



Lane 1 – MW markers; Lane 2 – ab84002; Lane 3 – ab84002 treated with PNGase F to remove potential N-linked glycans; Lane 4 – ab84002 treated with a glycosidase cocktail to remove potential N and O-linked glycans. Approximately 5 µg of protein was loaded per lane; Gel was stained using Deep Purple™. Drop in MW after treatment with PNGase F indicates presence of N-linked glycans. A further drop in MW after treatment with the glycosidase cocktail indicates the presence of O-linked glycans. Additional bands in lane 3 and lane 4 are glycosidase enzymes.



Post-translational modifications result in protein heterogeneity. The densitometry scan demonstrates the purified human cell expressed protein exists in multiple glycoforms, which differ according to their level of post-translational modification.

The triangle indicates theoretical pI and MW of the protein.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors