# abcam

# Product datasheet

# Recombinant Human H-Ras (G12V) protein (Tagged) ab90627

# 1 References

描述

产品名称 重组人H-Ras (G12V)蛋白(Tagged)

纯**度** > 95 % SDS-PAGE.

表达系统 Escherichia coli

Accession P01112

**蛋白长度** Full length protein

无动物成分 No

性质 Recombinant

种属 Human

序列 MTEYKLVVVG AGGVGKSALT IQLIQNHFVD

EYDPTIEDSY RKQVVIDGET CLLDILDTAG QEEYSAMRDQ YMRTGEGFLC VFAINNTKSF EDIHQYREQI KRVKDSDDVP MVLVGNKCDL AARTVESRQA QDLARSYGIP YIETSAKTRQ

GVEDAFYTLV REIRQHKLRK LNPPDESGPG CMSCKCVLS

预**测分子量** 21 kDa **氨基酸** 1 to 189

修饰 mutated G12V

标签 GST tag N-Terminus

额外的序列信息 (NP\_005334)

技术指标

Our **Abpromise guarantee** covers the use of **ab90627** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应**用** SDS-PAGE

**Functional Studies** 

形式 Liquid

补充说明 The mutation G12V leads to elimination of the intrinsic GTPase activity.

ab90627 is effective in activation of PI3K and PKB.

1

#### 制备和贮存

#### 稳定性和存储

Shipped on dry ice. Upon delivery aliquot and store at -80°C. Avoid freeze / thaw cycles.

pH: 7.50

 $Constituents: 0.87\% \ Sodium \ chloride, 0.395\% \ Tris \ HCl, 0.039\% \ Beta \ mercaptoethanol, 0.0475\% \ Annual Markov \ Markov$ 

Magnesium chloride

#### 常规信息

#### 功能

Ras proteins bind GDP/GTP and possess intrinsic GTPase activity.

#### 组织特异性

Widely expressed.

# 疾病相关

Defects in HRAS are the cause of faciocutaneoskeletal syndrome (FCSS) [MIM:218040]. A rare condition characterized by prenatally increased growth, postnatal growth deficiency, mental retardation, distinctive facial appearance, cardiovascular abnormalities (typically pulmonic stenosis, hypertrophic cardiomyopathy and/or atrial tachycardia), tumor predisposition, skin and musculoskeletal abnormalities.

Defects in HRAS are the cause of congenital myopathy with excess of muscle spindles (CMEMS) [MIM:218040]. CMEMS is a variant of Costello syndrome.

Defects in HRAS may be a cause of susceptibility to Hurthle cell thyroid carcinoma (HCTC) [MIM:607464]. Hurthle cell thyroid carcinoma accounts for approximately 3% of all thyroid cancers. Although they are classified as variants of follicular neoplasms, they are more often multifocal and somewhat more aggressive and are less likely to take up iodine than are other follicular neoplasms.

Note=Mutations which change positions 12, 13 or 61 activate the potential of HRAS to transform cultured cells and are implicated in a variety of human tumors.

Defects in HRAS are a cause of susceptibility to bladder cancer (BLC) [MIM:109800]. A malignancy originating in tissues of the urinary bladder. It often presents with multiple tumors appearing at different times and at different sites in the bladder. Most bladder cancers are transitional cell carcinomas. They begin in cells that normally make up the inner lining of the bladder. Other types of bladder cancer include squamous cell carcinoma (cancer that begins in thin, flat cells) and adenocarcinoma (cancer that begins in cells that make and release mucus and other fluids). Bladder cancer is a complex disorder with both genetic and environmental influences.

Note=Defects in HRAS are the cause of oral squamous cell carcinoma (OSCC).

#### 序列相似性

Belongs to the small GTPase superfamily. Ras family.

#### 翻译后修饰

Palmitoylated by the ZDHHC9-GOLGA7 complex. A continuous cycle of de- and re-palmitoylation regulates rapid exchange between plasma membrane and Golgi.

S-nitrosylated; critical for redox regulation. Important for stimulating guanine nucleotide exchange. No structural perturbation on nitrosylation.

The covalent modification of cysteine by 15-deoxy-Delta12,14-prostaglandin-J2 is autocatalytic and reversible. It may occur as an alternative to other cysteine modifications, such as S-nitrosylation and S-palmitoylation.

#### 细胞定位

Cell membrane. Cell membrane. Golgi apparatus. Golgi apparatus membrane. The active GTP-bound form is localized most strongly to membranes than the inactive GDP-bound form (By similarity). Shuttles between the plasma membrane and the Golgi apparatus and Nucleus. Cytoplasm. Cytoplasm > perinuclear region. Colocalizes with GNB2L1 to the perinuclear region.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

# Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.cn/abpromise">https://www.abcam.cn/abpromise</a> or contact our technical team.

#### Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors